(1/6)

Place of Sampling	Name of Sample (Region)		Analysis Item		
		Date of Sampling	Cs-134	Cs-137	Cs (Sum)
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))
Around 1km Offshore of Ota River (T-S1)	Blue crab (whole)	2022/7/22	< 3.5E+00	< 3.4E+00	ND
Around 1km Offshore of Ota River (T-S1)	Common skete (muscle)	2022/7/22	< 2.7E+00	< 3.8E+00	ND
Around 1km Offshore of Ota River (T-S1)	Flatfish (muscle) No.1	2022/7/22	< 4.9E+00	< 5.8E+00	ND
Around 1km Offshore of Ota River (T-S1)	Flatfish (muscle) No.2	2022/7/22	< 4.2E+00	< 2.8E+00	ND
Around 1km Offshore of Ota River (T-S1)	Marbled sole (muscle)	2022/7/22	< 3.8E+00	< 4.2E+00	ND
Around 3km Offshore of Odaka Ward (T-S2)	Lepidotrigla microptena (muscle)	2022/7/22	< 3.5E+00	< 3.3E+00	ND
Around 3km Offshore of Odaka Ward (T-S2)	Flatfish (muscle) No.1	2022/7/22	< 7.0E+00	< 5.8E+00	ND
Around 3km Offshore of Odaka Ward (T-S2)	Searobin (muscle)	2022/7/22	< 3.6E+00	< 3.6E+00	ND
Around 3km Offshore of Odaka Ward (T-S2)	Chub mackerel (muscle)	2022/7/22	< 3.1E+00	< 3.4E+00	ND
Around 3km Offshore of Ukedo River (T-S3)	Japanese angel shark (muscle)	2022/7/12	< 3.9E+00	< 3.9E+00	ND

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012): Sum of radioactivity concentrations for Cs-134 and Cs-137: 1.0E+02Bq/kg.
- · Analysis was conducted by Tokyo Power Technology Ltd.
- · Values are expressed in exponential notation. For example, "3.1E+01" means " 3.1×10^{1} " and equals 31. Similarly, "3.1E+00" means " 3.1×10^{0} " and equals 3.1, and "3.1E-01" means " 3.1×10^{-1} " and equals 0.31.

(2/6)

Place of Sampling	Name of Sample (Region)		Analysis Item		
		Date of Sampling	Cs-134	Cs-137	Cs (Sum)
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))
Around 3km Offshore of Ukedo River (T-S3)	Common skete (muscle)	2022/7/12	< 3.6E+00	< 3.5E+00	ND
Around 3km Offshore of Ukedo River (T-S3)	Flatfish (muscle) No.1	2022/7/12	< 6.1E+00	< 5.8E+00	ND
Around 3km Offshore of Ukedo River (T-S3)	Flatfish (muscle) No.2	2022/7/12	< 3.4E+00	< 3.2E+00	ND
Around 3km Offshore of Ukedo River (T-S3)	Searobin (muscle)	2022/7/12	< 3.5E+00	< 3.8E+00	ND
Around 3km Offshore of Ukedo River (T-S3)	Flathead (muscle)	2022/7/12	< 3.6E+00	< 3.4E+00	ND
Around 3km Offshore of 1F Site (T-S4)	Spiny dogfish (muscle)	2022/7/12	< 4.1E+00	< 3.6E+00	ND
Around 3km Offshore of 1F Site (T-S4)	Stone flounder (muscle)	2022/7/12	< 3.4E+00	3.1E+00	3.1E+00
Around 3km Offshore of 1F Site (T-S4)	Common skete (muscle)	2022/7/12	< 3.6E+00	5.8E+00	5.8E+00
Around 3km Offshore of 1F Site (T-S4)	Microstomus achne (muscle)	2022/7/12	< 3.6E+00	< 3.4E+00	ND
Around 3km Offshore of 1F Site (T-S4)	Flatfish (muscle) No.1	2022/7/12	< 5.2E+00	< 6.0E+00	ND

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012): Sum of radioactivity concentrations for Cs-134 and Cs-137: 1.0E+02Bq/kg.
- · Analysis was conducted by Tokyo Power Technology Ltd.
- · Values are expressed in exponential notation. For example, "3.1E+01" means " 3.1×10^{1} " and equals 31. Similarly, "3.1E+00" means " 3.1×10^{0} " and equals 3.1, and "3.1E-01" means " 3.1×10^{-1} " and equals 0.31.

(3/6)

			Analysis Item		
Place of Sampling	Name of Sample (Region)	Date of Sampling	Cs-134	Cs-137	Cs (Sum)
	(3 ,		(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))
Around 3km Offshore of 1F Site (T-S4)	Flatfish (muscle) No.2	2022/7/12	< 3.7E+00	< 4.0E+00	ND
Around 3km Offshore of 1F Site (T-S4)	Searobin (muscle)	2022/7/12	< 4.3E+00	< 3.0E+00	ND
Around 3km Offshore of 1F Site (T-S4)	Red sea bream (muscle)	2022/7/12	< 3.2E+00	< 3.7E+00	ND
Around 3km Offshore of 1F Site (T-S4)	John dory (muscle)	2022/7/12	< 3.1E+00	< 3.4E+00	ND
Around 2km Offshore of Kido River (T-S5)	Japanese angel shark (muscle)	2022/7/21	< 3.8E+00	< 3.9E+00	ND
Around 2km Offshore of Kido River (T-S5)	Common skete (muscle)	2022/7/21	< 4.2E+00	< 3.8E+00	ND
Around 2km Offshore of Kido River (T-S5)	Flatfish (muscle) No.1	2022/7/21	< 5.1E+00	< 5.0E+00	ND
Around 2km Offshore of Kido River (T-S5)	Flatfish (muscle) No.2	2022/7/21	< 3.7E+00	< 3.5E+00	ND
Around 2km Offshore of Kido River (T-S5)	Searobin (muscle)	2022/7/21	< 3.6E+00	< 3.0E+00	ND
Around 2km Offshore of 2F Site (T-S7)	Common skete (muscle)	2022/7/21	< 3.8E+00	< 3.5E+00	ND

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- \cdot Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012): Sum of radioactivity concentrations for Cs-134 and Cs-137: 1.0E+02Bq/kg.
- · Analysis was conducted by Tokyo Power Technology Ltd.
- · Values are expressed in exponential notation. For example, "3.1E+01" means " 3.1×10^{1} " and equals 31. Similarly, "3.1E+00" means " 3.1×10^{0} " and equals 3.1, and "3.1E-01" means " 3.1×10^{-1} " and equals 0.31.

(4/6)

Place of Sampling			Analysis Item			
	Name of Sample (Region)	Date of Sampling	Cs-134	Cs-137	Cs (Sum)	
	, ,		(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 2km Offshore of 2F Site (T-S7)	Crimson sea bream (muscle)	2022/7/21	< 4.3E+00	< 4.0E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Microstomus achne (muscle)	2022/7/21	< 3.4E+00	3.4E+00	3.4E+00	
Around 2km Offshore of 2F Site (T-S7)	Flatfish (muscle) No.1	2022/7/21	< 3.8E+00	< 3.1E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Flatfish (muscle) No.2	2022/7/21	< 3.5E+00	< 3.4E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Searobin (muscle)	2022/7/21	< 4.0E+00	< 3.7E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Red sea bream (muscle)	2022/7/21	< 3.3E+00	< 3.7E+00	ND	
Around 4km Offshore of Kumagawa (T-S8)	Japanese angel shark (muscle)	2022/7/14	< 3.6E+00	< 3.1E+00	ND	
Around 4km Offshore of Kumagawa (T-S8)	Lepidotrigla microptena (muscle)	2022/7/14	< 3.4E+00	< 3.4E+00	ND	
Around 4km Offshore of Kumagawa (T-S8)	Blue crab (whole)	2022/7/14	< 3.8E+00	< 3.7E+00	ND	
Around 4km Offshore of Kumagawa (T-S8)	Common skete (muscle)	2022/7/14	< 3.4E+00	< 3.6E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012): Sum of radioactivity concentrations for Cs-134 and Cs-137: 1.0E+02Bq/kg.
- · Analysis was conducted by Tokyo Power Technology Ltd.
- · Values are expressed in exponential notation. For example, "3.1E+01" means " 3.1×10^{1} " and equals 31. Similarly, "3.1E+00" means " 3.1×10^{0} " and equals 3.1, and "3.1E-01" means " 3.1×10^{-1} " and equals 0.31.

(5/6)

Place of Sampling			Analysis Item		
	Name of Sample (Region)	Date of Sampling	Cs-134	Cs-137	Cs (Sum)
	(3 /		(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))
Around 4km Offshore of Kumagawa (T-S8)	Flatfish (muscle) No.1	2022/7/14	< 3.3E+00	< 3.5E+00	ND
Around 4km Offshore of Kumagawa (T-S8)	Flatfish (muscle) No.2	2022/7/14	< 3.3E+00	< 3.6E+00	ND
Around 4km Offshore of Kumagawa (T-S8)	Searobin (muscle)	2022/7/14	< 2.9E+00	< 3.2E+00	ND
Around 15km Offshore of Odaka Ward (T-B1)	Mirror dory (muscle)	2022/8/5	< 3.6E+00	< 3.8E+00	ND
Around 15km Offshore of Odaka Ward (T-B1)	Lepidotrigla microptena (muscle)	2022/8/5	< 2.7E+00	< 3.8E+00	ND
Around 15km Offshore of Odaka Ward (T-B1)	Yellow goosefish (whole)	2022/8/5	< 3.9E+00	< 3.7E+00	ND
Around 15km Offshore of Odaka Ward (T-B1)	Common skete (muscle)	2022/8/5	< 3.9E+00	< 3.6E+00	ND
Around 15km Offshore of Odaka Ward (T-B1)	Crimson sea bream (muscle)	2022/8/5	< 3.8E+00	< 4.1E+00	ND
Around 15km Offshore of Odaka Ward (T-B1)	John dory (muscle)	2022/8/5	< 3.3E+00	< 2.9E+00	ND
Around 18km Offshore of Ukedo River (T-B2)	Mirror dory (muscle)	2022/8/5	< 4.3E+00	< 3.5E+00	ND

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012): Sum of radioactivity concentrations for Cs-134 and Cs-137: 1.0E+02Bq/kg.
- · Analysis was conducted by Tokyo Power Technology Ltd.
- · Values are expressed in exponential notation. For example, "3.1E+01" means " 3.1×10^{1} " and equals 31. Similarly, "3.1E+00" means " 3.1×10^{0} " and equals 3.1, and "3.1E-01" means " 3.1×10^{-1} " and equals 0.31.

(6/6)

			Analysis Item		
Place of Sampling	Name of Sample (Region)	Date of Sampling	Cs-134	Cs-137	Cs (Sum)
	(3, 7)		(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))
Around 18km Offshore of Ukedo River (T-B2)	Yellow goosefish (whole)	2022/8/5	< 4.2E+00	< 3.9E+00	ND
Around 18km Offshore of Ukedo River (T-B2)	Common skete (muscle)	2022/8/5	< 4.0E+00	< 3.4E+00	ND
Around 18km Offshore of Ukedo River (T-B2)	White croaker (muscle)	2022/8/5	< 4.0E+00	< 3.7E+00	ND
Around 18km Offshore of Ukedo River (T-B2)	Crimson sea bream (muscle)	2022/8/5	< 4.2E+00	< 4.0E+00	ND
Around 18km Offshore of Ukedo River (T-B2)	Red sea bream (muscle)	2022/8/5	< 4.1E+00	< 4.0E+00	ND
Around 18km Offshore of Ukedo River (T-B2)	John dory (muscle)	2022/8/5	< 4.0E+00	< 3.9E+00	ND

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- \cdot Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012): Sum of radioactivity concentrations for Cs-134 and Cs-137: 1.0E+02Bq/kg.
- · Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " 3.1×10^{1} " and equals 31. Similarly, "3.1E+00" means " 3.1×10^{0} " and equals 3.1, and "3.1E-01" means " 3.1×10^{1} " and equals 0.31.