

Evaluation of the exposure dose of workers engaged in radiation work at
the Fukushima Daiichi Nuclear Power Station

May 31, 2024

Tokyo Electric Power Company Holdings, Inc.
Fukushima Daiichi D & D Engineering Company

TEPCO has been evaluating the exposure dose of workers who engaged in radiation work at the Fukushima Daiichi Nuclear Power Station under two types, internal and external exposure to radiation, and has submitted the evaluation results to the Ministry of Health, Labour and Welfare regularly.

TEPCO today submitted to the Ministry of Health, Labour and Welfare a report on the exposure dose evaluation the data of which are those we collected until the end of April 2024. Here is part of the report: the maximum value of the external exposure dose among the workers who engaged in the work at the power station in April was 8.90 mSv, and regarding the internal exposure dose, no significant value was measured.

Exposure Dose Distribution

1. Effective Dose from External Exposure

Table 1 shows the distribution of external exposure dose of workers who were involved in radiation work at the Fukushima Daiichi Nuclear Power Station for the past three months.

Table 1. External Exposure Dose

Dose Ranges (mSv)	February 2024			March 2024			April 2024		
	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total
Above 100	0	0	0	0	0	0	0	0	0
75-100	0	0	0	0	0	0	0	0	0
50-75	0	0	0	0	0	0	0	0	0
20-50	0	0	0	0	0	0	0	0	0
10-20	0	4	4	0	0	0	0	0	0
5-10	0	35	35	0	37	37	0	27	27
1-5	3	468	471	12	440	452	12	367	379
1 or less	1041	6460	7501	1005	6422	7427	1003	6031	7034
Total	1044	6967	8011	1017	6899	7916	1015	6425	7440
Maximum (mSv)	1.9	10.8	10.8	3.9	8.6	8.6	3.22	8.90	8.90
Average (mSv)	0.06	0.30	0.26	0.08	0.28	0.26	0.07	0.24	0.22

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

2. Sum of External and Internal Exposure Dose (Effective Dose)

Table 2 shows the distribution of cumulative exposure dose of workers who are involved in radiation work at Fukushima Daiichi for five years, starting on April 1, 2021. Table 3 shows the distribution of cumulative exposure dose in the fiscal year of 2024. Two different periods of time are shown in the Table 2: from April 1, 2021 to March 31, 2024 and from April 1, 2021 to April 30, 2024. In the Table 3 the period from April 1, 2024 to April 30, 2024 is shown.

Table 2. Cumulative Exposure Dose for Five Years

Dose Ranges (mSv)	April 2021 - March 2024			April 2021 - April 2024			Difference		
	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total
Above 100	0	0	0	0	0	0	0	0	0
75-100	0	0	0	0	0	0	0	0	0
50-75	0	1	1	0	2	2	0	1	1
20-50	22	993	1015	25	1028	1053	3	35	38
10-20	55	1689	1744	54	1714	1768	-1	25	24
5-10	106	1576	1682	110	1585	1695	4	9	13
1-5	358	2647	3005	367	2657	3024	9	10	19
1 or less	1212	7840	9052	1201	7962	9163	-11	122	111
Total	1753	14746	16499	1757	14948	16705	4	202	206
Maximum (mSv)	31.12	50.82	50.82	31.36	51.88	51.88	-	-	-
Average (mSv)	1.79	4.94	4.60	1.83	4.97	4.64	-	-	-

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

• No significant internal exposure has been reported since October 2011.

Table 3. Cumulative Exposure Dose in the Fiscal Year of 2024

Dose Ranges (mSv)	April 2024		
	TEPCO Employees	Contractors	Total
Above 100	0	0	0
75-100	0	0	0
50-75	0	0	0
20-50	0	0	0
10-20	0	0	0
5-10	0	27	27
1-5	12	367	379
1 or less	1003	6031	7034
Total	1015	6425	7440
Maximum (mSv)	3.22	8.90	8.90
Average (mSv)	0.07	0.24	0.22

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

3. Sum of External and Internal Exposure Dose of Workers Exposed to Especially High Radiation (Effective Dose)

Table 4 shows the distribution of cumulative exposure dose of workers exposed to especially high radiation.*1

Table 4. Cumulative Exposure Dose (workers exposed to especially high radiation)

Dose Ranges (mSv)	March 2011 - September 2015
Above 100	1
75-100	191
50-75	233
20-50	267
10-20	186
5-10	129
1-5	145
1 or less	51
Total	1203
Maximum (mSv)	102.69
Average (mSv)	36.49

(Since October 2015, TEPCO Holdings has opted not to report to the Labour Standards Inspection Office about workers exposed to especially high radiation.)

*1. Workers exposed to especially high radiation means workers who are involved in operations in which they could be exposed to the emergency exposure dose limit (100 mSv), which is stipulated in "Ordinance on Prevention of Ionizing Radiation Hazards, Chapter 7." In more detail, they are workers engaged in the work to maintain the function of the cooling facility to cool down the reactor facility or the spent fuel tank in the reactor facility, the steam turbine and its related facilities or the surrounding area where the radiation doses exceed 0.1 mSv/h. Or they are workers who would engage in keeping running the function to control or prevent the release of a large number of radioactive materials should it be likely to occur due to malfunction or damage of the reactor facility.

So far workers who have worked as “workers exposed to especially high radiation” are all TEPCO employees.

*2. The figures in the cumulative data during the period from March 2011 to September 2015 in Table 4 above include the numbers of workers who have been reported to work as “workers exposed to especially high radiation” at least once.

*3. The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or the dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

*4. The figure shown in the dose range, “Above 100mSv,” in the cumulative data during the period from March 2011 to September 2015 is the figure when the March 2011 data of the internal exposure dose were reevaluated in July 2013.

4. Equivalent Dose

Table 5 and Table 6 show equivalent dose to the skin and the lens of the eye of the workers, respectively, who were involved in radiation work at the Fukushima Daiichi Nuclear Power Station for the past three months.

Table 5. Equivalent Dose to the Skin

Dose Ranges (mSv)	February 2024			March 2024			April 2024		
	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total
Above 500	0	0	0	0	0	0	0	0	0
300-500	0	0	0	0	0	0	0	0	0
250-300	0	0	0	0	0	0	0	0	0
200-250	0	0	0	0	0	0	0	0	0
150-200	0	0	0	0	0	0	0	0	0
100-150	0	0	0	0	0	0	0	0	0
75-100	0	0	0	0	0	0	0	0	0
50-75	0	0	0	0	0	0	0	0	0
20-50	0	0	0	0	1	1	0	0	0
10-20	0	4	4	0	1	1	0	0	0
5-10	0	59	59	0	59	59	0	27	27
1-5	6	545	551	19	546	565	12	422	434
1 or less	1038	6359	7397	998	6292	7290	1003	5976	6979
Total	1044	6967	8011	1017	6899	7916	1015	6425	7440
Maximum (mSv)	2.2	10.8	10.8	4.8	21.4	21.4	3.22	8.90	8.90
Average (mSv)	0.06	0.35	0.31	0.10	0.35	0.32	0.08	0.26	0.24

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or the dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

• Equivalent dose is a measure of the radiation dose to organs and tissues, and the equivalent dose limit to the skin is 500 mSv/year (the emergency exposure dose limit is 1 Sv).

• Equivalent dose to the skin is measured at a depth of 70 micrometers from the skin surface. When the equivalent dose is measured with a dosimeter other than the one put on around the chest and the abdomen, for example, a finger dosimeter, and the maximum measurement value is counted as the equivalent dose.

Table 6. Equivalent Dose to the Lens of the Eye

Dose Ranges (mSv)	February 2024			March 2024			April 2024		
	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total
Above 150	0	0	0	0	0	0	0	0	0
100-150	0	0	0	0	0	0	0	0	0
75-100	0	0	0	0	0	0	0	0	0
50-75	0	0	0	0	0	0	0	0	0
20-50	0	0	0	0	0	0	0	0	0
10-20	0	8	8	0	0	0	0	0	0
5-10	0	42	42	0	51	51	0	27	27
1-5	4	484	488	15	461	476	12	422	434
1 or less	1040	6433	7473	1002	6387	7389	1003	5976	6979
Total	1044	6967	8011	1017	6899	7916	1015	6425	7440
Maximum (mSv)	2.1	12.1	12.1	3.8	9.1	9.1	3.22	8.90	8.90
Average (mSv)	0.06	0.31	0.28	0.09	0.30	0.27	0.08	0.26	0.24

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or the dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

• Equivalent dose is a measure of the radiation dose to organs and tissues, and the equivalent dose limit to the lens of the eye is 50 mSv/year and 100 mSv/5 years (the emergency exposure dose limit is 300 mSv). The equivalent dose limit to the lens of the eye before April 1, 2021 was 150mSv/year (the emergency exposure dose limit was 300 mSv).

• The equivalent dose to the lens of the eye is measured at a depth of 1 centimeter for neutron ray, 3 millimeters for X-ray, gamma ray and beta ray from the skin surface. However, as for X-ray, gamma ray and beta ray, it is measured at a depth of 1 centimeter or 70 micrometer when deemed appropriate with consideration for radiation type and energy type (since April, 2021).

5. Cumulative Equivalent Dose

Table 7 and Table 8 show the distribution of cumulative equivalent dose to the skins and the lens of the eye of the workers, respectively, who were involved in radiation work at the Fukushima Daiichi Nuclear Power Station from April 1, 2024 to April 30, 2024.

Table 9 shows the distribution of cumulative exposure dose for five years, starting on April 1, 2021: from April 1, 2021 to March 31, 2024 and from April 1, 2021 to April 30, 2024 for comparison.

Table 7. Equivalent Dose to the Skin

Dose Ranges (mSv)	April 2024		
	TEPCO Employees	Contractors	Total
Above 500	0	0	0
300-500	0	0	0
250-300	0	0	0
200-250	0	0	0
150-200	0	0	0
100-150	0	0	0
75-100	0	0	0
50-75	0	0	0
20-50	0	0	0
10-20	0	0	0
5-10	0	27	27
1-5	12	422	434
1 or less	1003	5976	6979
Total	1015	6425	7440
Maximum (mSv)	3.22	8.90	8.90
Average (mSv)	0.08	0.26	0.24

- The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or the dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

- Equivalent dose is a measure of the radiation dose to organs and tissues, and the equivalent dose limit to the skin is 500 mSv/year (the emergency exposure dose limit is 1 Sv).

- Equivalent dose to the skin is measured at a depth of 70 micrometers from the skin surface. When the equivalent dose is measured with a dosimeter other than the one put on around the chest and the abdomen, for example, a finger dosimeter, and the maximum measurement value is counted as the equivalent dose.

Table 8. Equivalent Dose to the Lens of the Eye

Dose Ranges (mSv)	April 2024		
	TEPCO Employees	Contractors	Total
Above 150	0	0	0
100-150	0	0	0
75-100	0	0	0
50-75	0	0	0
20-50	0	0	0
10-20	0	0	0
5-10	0	27	27
1-5	12	422	434
1 or less	1003	5976	6979
Total	1015	6425	7440
Maximum (mSv)	3.22	8.90	8.90
Average (mSv)	0.08	0.26	0.24

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or the dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

• Equivalent dose is a measure of the radiation dose to organs and tissues, and the equivalent dose limit to the lens of the eye is 50 mSv/year and 100 mSv/5 years (the emergency exposure dose limit is 300 mSv).

• The equivalent dose to the lens of the eye is measured at a depth of 1 centimeter for neutron ray, 3 millimeters for X-ray, gamma ray and beta ray from the skin surface. However, as for X-ray, gamma ray and beta ray, it is measured at a depth of 1 centimeter or 70 micrometer when deemed appropriate with consideration for radiation type and energy type.

Table 9. Equivalent Dose to the Lens of the Eye: Cumulative Exposure Dose for Five Years

Dose Ranges (mSv)	April 2021 - March 2024			April 2021 - April 2024			Difference		
	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total	TEPCO Employees	Contractors	Total
Above 100	0	0	0	0	0	0	0	0	0
75-100	0	0	0	0	0	0	0	0	0
50-75	0	0	0	0	6	6	0	6	6
20-50	24	1097	1121	26	1133	1159	2	36	38
10-20	55	1683	1738	55	1723	1778	0	40	40
5-10	110	1529	1639	114	1510	1624	4	-19	-15
1-5	359	2631	2990	370	2650	3020	11	19	30
1 or less	1205	7806	9011	1192	7926	9118	-13	120	107
Total	1753	14746	16499	1757	14948	16705	4	202	206
Maximum (mSv)	31.31	50.00	50.00	31.55	51.13	51.13	-	-	-
Average (mSv)	1.81	5.15	4.79	1.85	5.19	4.84	-	-	-

• The values of the exposure dose and the number of the workers in the table above are subject to change, because there are cases that APD data are replaced with monthly dose data measured by integral dosimeters. Or the dose data of workers who wore only an integral dosimeter (ex., workers who entered only the Seismic Isolation Building) need to be updated in the table after the publication of the data.

• Equivalent dose is a measure of the radiation dose to organs and tissues, and the equivalent dose limit to the lens of the eye is 50 mSv/year and 100 mSv/5 years (the emergency exposure dose limit is 300 mSv).

• The equivalent dose to the lens of the eye is measured at a depth of 1 centimeter for neutron ray, 3 millimeters for X-ray, gamma ray and beta ray from the skin surface. However, as for X-ray, gamma ray and beta ray, it is measured at a depth of 1 centimeter or 70 micrometer when deemed appropriate with consideration for radiation type and energy type.