Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station> <1/2>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle) No.1	Port area of Fukushima Daiichi NPS (Shallow draft quay)	February 25, 2020	ND(2.5)	9.5	9.5
Marbled sole (muscle) No.2	Port area of Fukushima Daiichi NPS (North breakwater)	February 25, 2020	ND(3.7)	43	43
Marbled sole (muscle) No.3	Port area of Fukushima Daiichi NPS (North breakwater)	February 25, 2020	ND(3.0)	5.0	5.0
Marbled sole (muscle) No.4	Port area of Fukushima Daiichi NPS (North side of the east wave breaker)	February 10, 2020	4.3	84	88.3
Roundnose flounder (muscle)	Port area of Fukushima Daiichi NPS (North side of the east wave breaker)	February 10, 2020	ND(2.1)	2.6	2.6
Common skete (muscle) No.1	Port area of Fukushima Daiichi NPS (North side of the east wave breaker)	February 20, 2020	ND(2.4)	57	57
Common skete (muscle) No.2	Port area of Fukushima Daiichi NPS (North side of the east wave breaker)	February 25, 2020	ND(2.5)	36	36
Flatfish (muscle)	Port area of Fukushima Daiichi NPS (South side of the east wave breaker)	February 5, 2020	9.7	150	159.7
Marbled sole (muscle) No.5	Port area of Fukushima Daiichi NPS (South side of the east wave breaker)	February 10, 2020	ND(2.8)	11	11
Marbled sole (muscle) No.6	Port area of Fukushima Daiichi NPS (South side of the east wave breaker)	February 10, 2020	ND(2.4)	4.3	4.3

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station> <2/2>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)			
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)	
Marbled sole (muscle) No.7	Port area of Fukushima Daiichi NPS (South side of the east wave breaker)	February 10, 2020	ND(2.6)	ND(3.0)	ND	
Common skete (muscle) No.3	Port area of Fukushima Daiichi NPS (South side of the east wave breaker)	February 20, 2020	ND(2.9)	22	22	
Marbled sole (muscle) No.8	Port area of Fukushima Daiichi NPS (South breakwater)	February 10, 2020	ND(2.8)	4.9	4.9	

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.