

Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle) No.1	Port area of Fukushima Daiichi NPS (Around the middle of port)	April 22, 2020	ND(2.6)	25	25
Marbled sole (muscle) No.2	Port area of Fukushima Daiichi NPS (Near shallow draft quay)	April 6, 2020	ND(1.8)	8.0	8.0
Marbled sole (muscle) No.3	Port area of Fukushima Daiichi NPS (Near shallow draft quay)	April 6, 2020	ND(2.9)	4.9	4.9
Marbled sole (muscle) No.4	Port area of Fukushima Daiichi NPS (Near north breakwater)	April 22, 2020	ND(2.2)	12	12
Marbled sole (muscle) No.5	Port area of Fukushima Daiichi NPS (Near south breakwater)	April 6, 2020	ND(2.1)	9.3	9.3

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.