

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <1/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Lepidotrigla microptena (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(3.5) | ND(3.6) | ND |
| Black rockfish (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(3.7) | ND(3.4) | ND |
| Black sea bream (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(3.4) | ND(3.1) | ND |
| Common skete (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(4.1) | ND(3.8) | ND |
| Microstomus achne (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(4.0) | ND(3.2) | ND |
| Flatfish (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(4.0) | ND(3.6) | ND |
| Marbled sole (muscle) | Around 1km Offshore of Ota River (T-S1) | April 10, 2020 | ND(3.3) | ND(3.5) | ND |
| Stone flounder (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(3.4) | ND(3.5) | ND |
| Lepidotrigla microptena (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(4.0) | ND(3.9) | ND |
| Black rockfish (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(3.9) | ND(2.9) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <2/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|-------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Common skete (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(3.9) | ND(3.9) | ND |
| Sea bass (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(4.6) | ND(3.8) | ND |
| Microstomus achne (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(3.1) | ND(3.5) | ND |
| Flatfish (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(2.5) | ND(3.7) | ND |
| Littlemouth flounder (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(3.3) | ND(3.4) | ND |
| Marbled sole (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(3.2) | ND(3.7) | ND |
| Roundnose flounder (muscle) | Around 3km Offshore of Odaka Ward (T-S2) | April 10, 2020 | ND(4.0) | ND(4.3) | ND |
| Stone flounder (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(3.9) | ND(3.0) | ND |
| Lepidotrigla microptena (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(4.1) | ND(3.7) | ND |
| Common skete (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(3.7) | ND(3.5) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
 (Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <3/11>
 (excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|----------------------------------|-------------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Microstomus achne (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(3.1) | 9.3 | 9.3 |
| Flatfish (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(4.0) | ND(3.8) | ND |
| Littlemouth flounder (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(3.8) | ND(3.3) | ND |
| Marbled sole (muscle) | Around 3km Offshore of Ukedo River (T-S3) | April 23, 2020 | ND(3.7) | ND(3.2) | ND |
| Roundnose flounder (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.4) | ND(3.4) | ND |
| Stone flounder (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.3) | ND(3.8) | ND |
| Black rockfish (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.7) | ND(3.0) | ND |
| Common skete (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(4.2) | ND(3.9) | ND |
| Flatfish (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.5) | ND(2.8) | ND |
| Littlemouth flounder (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.3) | ND(4.1) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <4/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|-------------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Marbled sole (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.6) | ND(3.8) | ND |
| Roundnose flounder (muscle) | Around 3km Offshore of Fukushima Daiichi (T-S4) | April 23, 2020 | ND(3.4) | ND(3.6) | ND |
| Lepidotrigla microptena (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.7) | ND(4.4) | ND |
| Black rockfish (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.6) | ND(3.6) | ND |
| Common skate (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.8) | 3.7 | 3.7 |
| Microstomus achne (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.8) | ND(3.4) | ND |
| Flatfish (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.6) | ND(4.3) | ND |
| Smooth dogfish (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.4) | ND(3.5) | ND |
| Marbled sole (muscle) | Around 2km Offshore of Kido River (T-S5) | April 28, 2020 | ND(3.8) | 4.1 | 4.1 |
| Stone flounder (muscle) | Around 2km Offshore of Fukushima Daini (T-S7) | April 28, 2020 | ND(3.8) | ND(3.9) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <5/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|-----------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Japanese angel shark (muscle) | Around 2km Offshore of Fukushima Daini (T-S7) | April 28, 2020 | ND(3.7) | ND(3.6) | ND |
| Lepidotrigla microptena (muscle) | Around 2km Offshore of Fukushima Daini (T-S7) | April 28, 2020 | ND(3.2) | ND(3.2) | ND |
| Common skete (muscle) | Around 2km Offshore of Fukushima Daini (T-S7) | April 28, 2020 | ND(3.4) | ND(3.9) | ND |
| Flatfish (muscle) | Around 2km Offshore of Fukushima Daini (T-S7) | April 28, 2020 | ND(4.0) | ND(3.7) | ND |
| Stone flounder (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.3) | ND(3.6) | ND |
| Lepidotrigla microptena (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(4.3) | ND(4.2) | ND |
| Black sea bream (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.1) | ND(3.3) | ND |
| Common skete (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(4.3) | ND(4.2) | ND |
| Microstomus achne (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.6) | ND(3.4) | ND |
| Flatfish (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.6) | ND(3.8) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <6/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|----------------------------------|-------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Searobin (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.5) | ND(3.7) | ND |
| Smooth dogfish (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.8) | ND(3.3) | ND |
| Littlemouth flounder (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.0) | ND(3.4) | ND |
| Marbled sole (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.2) | ND(3.7) | ND |
| Flathead (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.9) | ND(3.9) | ND |
| Ridged-eye flounder (muscle) | Around 4km Offshore of Kumagawa (T-S8) | April 23, 2020 | ND(3.8) | ND(3.8) | ND |
| Stone flounder (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(4.3) | ND(3.5) | ND |
| Lepidotrigla microptena (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.9) | ND(3.7) | ND |
| Common skete (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.1) | ND(3.3) | ND |
| White croaker (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.4) | ND(3.3) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <7/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|--------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Sea bass (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.7) | ND(3.6) | ND |
| Pointhead flounder (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.5) | ND(3.9) | ND |
| Microstomus achne (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.2) | ND(3.9) | ND |
| Flatfish (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(2.8) | ND(3.3) | ND |
| Searobin (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(3.3) | ND(3.6) | ND |
| Marbled sole (muscle) | Around 15km Offshore of Odaka Ward (T-B1) | April 24, 2020 | ND(4.3) | ND(4.0) | ND |
| Greenling (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.6) | ND(4.2) | ND |
| Stone flounder (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(4.0) | ND(3.0) | ND |
| Lepidotrigla microptena (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.8) | ND(3.7) | ND |
| Yellow goosfish (whole) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.7) | ND(4.1) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <8/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|------------------------------------|--------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Common skate (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.8) | ND(3.8) | ND |
| White croaker (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.8) | ND(3.4) | ND |
| Sea bass (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(4.3) | 4.1 | 4.1 |
| Pointhead flounder (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(4.0) | ND(3.9) | ND |
| Cloudy catshark (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.2) | ND(3.6) | ND |
| Microstomus achne (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(4.3) | ND(3.0) | ND |
| Flatfish (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.3) | ND(3.7) | ND |
| Common Japanese conger (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.1) | ND(3.9) | ND |
| Littlemouth flounder (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.7) | ND(3.5) | ND |
| Marbled sole (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(4.0) | ND(3.0) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <9/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|--------------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Willow flounder (muscle) | Around 18km Offshore of Ukedo River (T-B2) | April 24, 2020 | ND(3.6) | ND(3.8) | ND |
| Greenling (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.4) | ND(3.3) | ND |
| Lepidotrigla microptena (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.5) | ND(3.6) | ND |
| Black rockfish (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.5) | ND(3.6) | ND |
| Common skate (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.5) | ND(2.9) | ND |
| Sea bass (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.5) | ND(4.3) | ND |
| Pointhead flounder (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(2.8) | ND(2.9) | ND |
| Crimson sea bream (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(4.1) | ND(4.1) | ND |
| Flatfish ① (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.6) | ND(4.0) | ND |
| Flatfish ② (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(3.3) | ND(4.2) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <10/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|-------------------------------------|--------------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Searobin (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(2.9) | ND(2.9) | ND |
| Littlemouth flounder (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(4.1) | ND(3.9) | ND |
| Marbled sole (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(4.0) | ND(4.5) | ND |
| Roundnose flounder (muscle) | Around 10km Offshore of Fukushima Daiichi (T-B3) | April 21, 2020 | ND(4.1) | ND(3.5) | ND |
| Greenling (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.7) | ND(3.1) | ND |
| Lepidotrigla microptena (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.8) | ND(2.9) | ND |
| Black rockfish (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.8) | ND(3.9) | ND |
| Common skate (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(4.1) | ND(4.1) | ND |
| Sea bass (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.6) | 3.9 | 3.9 |
| Pointhead flounder (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.3) | ND(3.6) | ND |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <11/11>
(excluding the port)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|---------------------------------|------------------------------------------------|------------------|-------------------------------------------------------|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Microstomus achne (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(4.2) | ND(3.9) | ND |
| Flatfish ① (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.4) | ND(4.6) | ND |
| Flatfish ② (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.4) | ND(3.7) | ND |
| Searobin (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.9) | ND(4.0) | ND |
| Marbled sole (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(4.3) | ND(4.0) | ND |
| Globefish (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.9) | ND(3.8) | ND |
| Roundnose flounder (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.2) | ND(3.7) | ND |
| Ridged-eye flounder (muscle) | Around 10km Offshore of Fukushima Daini (T-B4) | April 21, 2020 | ND(3.6) | ND(3.6) | ND |
| | | | | | |
| | | | | | |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

*Analysis was conducted by Tokyo Power Technology Ltd.