Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station> <1/1>

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|----------------------------|---|------------------|---|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Flatfish (muscle) No.1 | Port area of Fukushima Daiichi NPS (Near south breakwater) | June 2, 2020 | ND(3.0) | 9.6 | 9.6 |
| Flatfish (muscle) No.2 | Port area of Fukushima Daiichi NPS (Near port entrance) | June 26, 2020 | ND(2.3) | 8.7 | 8.7 |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.