(1/7)

Place of Sampling	Name of Sample (Region)	Date of Sampling	Analysis Item			
			Cs-134	Cs-137	Cs (Sum)	
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 1km Offshore of Ota River (T-S1)	Black rockfish (muscle)	2021/1/14	< 3.6E+00	< 4.0E+00	ND	
Around 1km Offshore of Ota River (T-S1)	Sea raven (muscle)	2021/1/14	< 3.7E+00	< 3.8E+00	ND	
Around 1km Offshore of Ota River (T-S1)	Common skete (muscle)	2021/1/14	< 4.3E+00	< 3.8E+00	ND	
Around 1km Offshore of Ota River (T-S1)	Marbled sole (muscle)	2021/1/14	< 3.4E+00	< 3.7E+00	ND	
Around 1km Offshore of Ota River (T-S1)	Roundnose flounder (muscle)	2021/1/14	< 3.3E+00	< 4.2E+00	ND	
Around 3km Offshore of Odaka Ward (T-S2)	Stone flounder (muscle)	2021/1/14	< 3.1E+00	< 3.7E+00	ND	
Around 3km Offshore of Odaka Ward (T-S2)	Black rockfish (muscle)	2021/1/14	< 3.8E+00	< 3.5E+00	ND	
Around 3km Offshore of Odaka Ward (T-S2)	Marbled sole (muscle)	2021/1/14	< 3.6E+00	< 3.8E+00	ND	
Around 3km Offshore of Odaka Ward (T-S2)	Roundnose flounder (muscle)	2021/1/14	< 3.3E+00	< 3.9E+00	ND	
Around 3km Offshore of Ukedo River (T-S3)	Stone flounder (muscle)	2021/1/15	< 3.7E+00	< 3.6E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- $\cdot \ \text{Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND)}.$
- $\cdot \ \text{Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.}$
- $\boldsymbol{\cdot}$  Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1 \times 10^{1}$ " and equals 31. Similarly, "3.1E+00" means " $3.1 \times 10^{0}$ " and equals 3.1, and " $3.1 \times 10^{0}$ " means " $3.1 \times 10^{0}$ " and equals 0.31.

(2/7)

Place of Sampling	Name of Sample (Region)	Date of Sampling	Analysis Item			
			Cs-134	Cs-137	Cs (Sum)	
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 3km Offshore of Ukedo River (T-S3)	Common skete (muscle)	2021/1/15	< 3.5E+00	< 4.3E+00	ND	
Around 3km Offshore of Ukedo River (T-S3)	Flatfish (muscle) No.1	2021/1/15	< 3.7E+00	< 3.6E+00	ND	
Around 3km Offshore of Ukedo River (T-S3)	Littlemouth flounder (muscle)	2021/1/15	< 4.1E+00	< 3.4E+00	ND	
Around 3km Offshore of Ukedo River (T-S3)	Marbled sole (muscle)	2021/1/15	< 3.7E+00	< 3.8E+00	ND	
Around 3km Offshore of Ukedo River (T-S3)	Flathead (muscle)	2021/1/15	< 3.7E+00	< 3.4E+00	ND	
Around 3km Offshore of Ukedo River (T-S3)	Roundnose flounder (muscle)	2021/1/15	< 4.1E+00	< 3.3E+00	ND	
Around 3km Offshore of 1F Site (T-S4)	Stone flounder (muscle)	2021/1/15	< 4.2E+00	< 3.7E+00	ND	
Around 3km Offshore of 1F Site (T-S4)	Japanese angel shark (muscle)	2021/1/15	< 3.8E+00	4.6E+00	4.6E+00	
Around 3km Offshore of 1F Site (T-S4)	Lepidotrigla microptena (muscle)	2021/1/15	< 3.6E+00	< 3.1E+00	ND	
Around 3km Offshore of 1F Site (T-S4)	Common skete (muscle)	2021/1/15	< 3.4E+00	< 3.5E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.
- Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1 \times 10^{1}$ " and equals 31. Similarly, "3.1E+00" means " $3.1 \times 10^{0}$ " and equals 3.1, and " $3.1 \times 10^{0}$ " means " $3.1 \times 10^{0}$ " and equals 0.31.

(3/7)

Place of Sampling	Name of Sample (Region)	Date of Sampling	Analysis Item			
			Cs-134	Cs-137	Cs (Sum)	
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 3km Offshore of 1F Site (T-S4)	Flatfish (muscle) No.1	2021/1/15	< 3.8E+00	< 3.3E+00	ND	
Around 3km Offshore of 1F Site (T-S4)	Marbled sole (muscle)	2021/1/15	< 3.8E+00	< 3.4E+00	ND	
Around 3km Offshore of 1F Site (T-S4)	Pacific cod (muscle)	2021/1/15	< 3.6E+00	< 3.8E+00	ND	
Around 3km Offshore of 1F Site (T-S4)	Roundnose flounder (muscle)	2021/1/15	< 3.9E+00	< 3.7E+00	ND	
Around 2km Offshore of Kido River (T-S5)	Stone flounder (muscle)	2021/1/22	< 3.5E+00	< 3.6E+00	ND	
Around 2km Offshore of Kido River (T-S5)	Japanese angel shark (muscle)	2021/1/22	< 3.7E+00	< 3.8E+00	ND	
Around 2km Offshore of Kido River (T-S5)	Common skete (muscle)	2021/1/22	< 3.6E+00	< 3.5E+00	ND	
Around 2km Offshore of Kido River (T-S5)	Microstomus achne (muscle)	2021/1/22	< 3.9E+00	< 3.3E+00	ND	
Around 2km Offshore of Kido River (T-S5)	Smooth dogfish (muscle)	2021/1/22	< 3.2E+00	5.6E+00	5.6E+00	
Around 2km Offshore of Kido River (T-S5)	Marbled sole (muscle)	2021/1/22	< 4.1E+00	< 3.9E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.
- Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1 \times 10^{1}$ " and equals 31. Similarly, "3.1E+00" means " $3.1 \times 10^{0}$ " and equals 3.1, and " $3.1 \times 10^{0}$ " means " $3.1 \times 10^{0}$ " and equals 0.31.

(4/7)

Place of Sampling	Name of Sample (Region)	Date of Sampling	Analysis Item			
			Cs-134	Cs-137	Cs (Sum)	
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 2km Offshore of Kido River (T-S5)	Roundnose flounder (muscle)	2021/1/22	< 2.9E+00	< 3.6E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Lepidotrigla microptena (muscle)	2021/1/22	< 3.8E+00	< 3.7E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Black rockfish (muscle)	2021/1/22	< 3.2E+00	3.9E+00	3.9E+00	
Around 2km Offshore of 2F Site (T-S7)	Common skete (muscle)	2021/1/22	< 3.7E+00	8.7E+00	8.7E+00	
Around 2km Offshore of 2F Site (T-S7)	Microstomus achne (muscle)	2021/1/22	< 3.7E+00	< 3.9E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Smooth dogfish (muscle)	2021/1/22	< 3.3E+00	< 3.8E+00	ND	
Around 2km Offshore of 2F Site (T-S7)	Roundnose flounder (muscle)	2021/1/22	< 4.1E+00	< 3.2E+00	ND	
Around 4km Offshore of Kumagawa (T-S8)	Common skete (muscle)	2021/1/21	< 3.4E+00	4.8E+00	4.8E+00	
Around 4km Offshore of Kumagawa (T-S8)	Microstomus achne (muscle)	2021/1/21	< 3.6E+00	< 3.8E+00	ND	
Around 4km Offshore of Kumagawa (T-S8)	Flatfish (muscle) No.1	2021/1/21	< 3.8E+00	< 3.4E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.
- $\boldsymbol{\cdot}$  Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1\times10^{11}$ " and equals 31. Similarly, "3.1E+00" means " $3.1\times10^{01}$ " and equals 3.1, and "3.1E-01" means " $3.1\times10^{-11}$ " and equals 0.31.

(5/7)

Place of Sampling	Name of Sample (Region)	Date of Sampling	Analysis Item			
			Cs-134	Cs-137	Cs (Sum)	
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 4km Offshore of Kumagawa (T-S8)	Roundnose flounder (muscle)	2021/1/21	< 3.3E+00	< 4.4E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Lepidotrigla microptena (muscle)	2021/1/13	< 2.9E+00	< 3.8E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Common skete (muscle)	2021/1/13	< 3.0E+00	< 3.6E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Takifugu snyderi (muscle)	2021/1/13	< 3.6E+00	< 3.4E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Sea bass (muscle)	2021/1/13	< 3.3E+00	< 3.1E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Flatfish (muscle) No.1	2021/1/13	< 4.5E+00	< 3.6E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Searobin (muscle)	2021/1/13	< 3.9E+00	< 3.4E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Smooth dogfish (muscle)	2021/1/13	< 3.3E+00	< 3.5E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Littlemouth flounder (muscle)	2021/1/13	< 4.2E+00	< 3.3E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Marbled sole (muscle)	2021/1/13	< 3.4E+00	< 3.8E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.
- $\boldsymbol{\cdot}$  Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1\times10^{11}$ " and equals 31. Similarly, "3.1E+00" means " $3.1\times10^{01}$ " and equals 3.1, and "3.1E-01" means " $3.1\times10^{-11}$ " and equals 0.31.

(6/7)

Place of Sampling	Name of Sample (Region)	Date of Sampling	Analysis Item			
			Cs-134	Cs-137	Cs (Sum)	
			(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 10km Offshore of 1F Site (T-B3)	Roundnose flounder (muscle)	2021/1/13	< 2.8E+00	< 3.8E+00	ND	
Around 10km Offshore of 1F Site (T-B3)	Ridged-eye flounder (muscle)	2021/1/13	< 3.4E+00	< 3.7E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Lepidotrigla microptena (muscle)	2021/1/13	< 3.1E+00	< 3.6E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Yellow goosefish (whole)	2021/1/13	< 3.3E+00	< 2.8E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Common skete (muscle)	2021/1/13	< 3.3E+00	< 3.6E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Takifugu snyderi (muscle)	2021/1/13	< 3.6E+00	< 3.7E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Sea bass (muscle)	2021/1/13	< 3.6E+00	< 2.6E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Microstomus achne (muscle)	2021/1/13	< 2.8E+00	< 3.4E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Flatfish (muscle) No.1	2021/1/13	< 4.1E+00	< 3.7E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Smooth dogfish (muscle)	2021/1/13	< 3.1E+00	< 4.2E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- · Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.
- $\boldsymbol{\cdot}$  Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1\times10^{11}$ " and equals 31. Similarly, "3.1E+00" means " $3.1\times10^{01}$ " and equals 3.1, and "3.1E-01" means " $3.1\times10^{-11}$ " and equals 0.31.

(7/7)

			Analysis Item			
Place of Sampling	Name of Sample (Region)	Date of Sampling	Cs-134	Cs-137	Cs (Sum)	
	(riagion)		(Bq/kg(Raw))	(Bq/kg(Raw))	(Bq/kg(Raw))	
Around 10km Offshore of 2F Site (T-B4)	Littlemouth flounder (muscle)	2021/1/13	< 3.1E+00	< 3.7E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Marbled sole (muscle)	2021/1/13	< 4.1E+00	< 3.9E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Roundnose flounder (muscle)	2021/1/13	< 3.2E+00	< 2.8E+00	ND	
Around 10km Offshore of 2F Site (T-B4)	Ridged-eye flounder (muscle)	2021/1/13	< 3.2E+00	< 3.2E+00	ND	

- · Half life of each nuclide: Cs-134 (Approx. 2 years), Cs-137 (Approx. 30 years)
- $\cdot$  Inequality sign (<: less than) indicates that measurement result is less than the detection limit (ND).
- $\cdot \ \text{Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.}$
- $\boldsymbol{\cdot}$  Analysis was conducted by Tokyo Power Technology Ltd.
- Values are expressed in exponential notation. For example, "3.1E+01" means " $3.1 \times 10^{1}$ " and equals 31. Similarly, "3.1E+00" means " $3.1 \times 10^{0}$ " and equals 3.1, and "3.1E-01" means " $3.1 \times 10^{-1}$ " and equals 0.31.