

Nuclide analysis results of ocean soil < 1/2 >

Reference

(Data summarized on March 3)

Place of Sampling	North of Discharge Channel of 5-6u of 1F (approx. 30m north of 5-6u discharge channel)	Around South Discharge Channel of 1F (1-4u Discharge Channel)	Around North Discharge Channel of 2F (Around 3,4u Discharge Channel) (approx. 10 km from 1F)	Around Iwasawa Shore of 2F (approx. 7 km south of 1,2u Discharge Channel) (approx. 16 km from 1F)	Iwasawa offshore 3km
Time of Sampling	Mar 01, 2012 9:10 AM	Mar 01, 2012 9:35 AM	Mar 01, 2012 (Not sampled)	Mar 01, 2012 8:05 AM	Mar 01, 2012 (Not sampled)
Detected Nuclides (Half-life)	Radioactivity density (Bq/kg· moist soil)				
I-131 (about 8 days)	ND	ND	-	ND	-
Cs-134 (about 2 years)	690	620	-	130	-
Cs-137 (about 30 years)	930	830	-	170	-

* Data of other nuclides are under evaluation.

* "ND" means the sampled data is below measurable limit.

I-131: approx. 8Bq/kg· moist soil)

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.


Nuclide analysis results of ocean soil < 2/2 >

Reference


(Data summarized on March 3)

Place of Sampling	8km offshore of Iwasawa shore				
Time of Sampling	Mar 01, 2012 (Not sampled)				
Detected Nuclides (Half-life)	Radioactivity density (Bq/kg· moist soil)				
I-131 (about 8 days)	-				
Cs-134 (about 2 years)	-				
Cs-137 (about 30 years)	-				

Radioactivity Density of Ocean Soil at North Discharge Channel of Fukushima Daiichi
NPS
(Around 5,6u Discharge Channel) (Bq/kg (moist soil))



Radioactivity Density of Ocean Soil at South Discharge Channel of Fukushima Daiichi NPS (Bq/kg (moist soil))



Radioactivity Density of Ocean Soil at Iwasawa Shore of Fukushima Daini NPS (Bq/kg (moist soil))

