Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station >

(Data summarized on November 1)

Place of Sampling	North of Unit 5-6 Discharge Daiichi N (Approx. 30m North of Unit 5	IPS	Around South Discharge C Daiichi N (Appox. 330m South of Unit	Density Limit Specified by the Reactor Regulation (Bq/L)		
Time of Sampling	Oct 31, 2 8:15 A		Oct 31, 2 8:00 A	(The density limit in the water outside the surrounding monitored areas is provided		
Detected Nuclides (Half-life)	Density of Sample (Bq/L)	Scaling Factor	Density of Sample (Bq/L)	Scaling Factor	in section 6 of Appendix 2.)	
I-131 (Approx. 8 days)	ND	-	ND	-	40	
Cs-134 (Approx. 2 years)	ND	-	ND	-	60	
Cs-137 (Approx. 30 years)	ND	-	ND	-	90	

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

I-131: Approx. 0.49Bq/L, Cs-134: Approx.1.1Bq/L, Cs-137: Approx.1.4Bq/L

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.

Nuclides Analysis Result of Radioactive Materials in the Seawater < Offshore >

(Data summarized on November 1)


Place of Sampling (Place No.)	Arounmd 15km Offshore of Odaka Ward (T-B1)				Around 18km Offshore of Ukedo River (T-B2)								Density Limit Specified by the Reactor Regulation
Upper Layer		Lower Layer		Upper Layer		Lower Layer		Upper Layer		Lower Layer		(Bq/L)	
Time of Sampling	September (Not sam		September (Not samp			September 2012 (Not sampled)		September 2012 (Not sampled)					(The density limit in the water outside the surrounding monitored
Detected Nuclides (Half-life)	Density of Sample (Bq/L)	Scaling Factor	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor	Density of Sample (Bq/L)	Scaling Factor	areas is provided in section 6 of Appendix 2.)
Cs-134 (Approx. 2 years)	-	-	-	-	-	-	-	-					60
Cs-137 (Approx. 30 years)	-	-	-	-	-	-	-	-					90

Place of Sampling (Place No.)	Upper La	Lower La	Upper Layer Lower Layer				Upper Layer Lower Layer				Density Limit Specified by the Reactor Regulation (Bq/L)		
Time of Sampling													(The density limit in the water outside the surrounding monitored
Detected Nuclides (Half-life)	Density of Sample (Bq/L)	Scaling Factor	Density of Sample (Bq/L)	Scaling Factor (/)	Density of Sample (Bq/L)	Scaling Factor (/)	aroos is provided in						
Cs-134 (Approx. 2 years)													60
Cs-137 (Approx. 30 years)													90

 $^{^{\}star}$ The density specified by the Reactor Regulation is converted from Bq/cm $^{\!3}$ to Bq/L.

No sampling due to the bad weather.

Radioactivity Density of the Seawater at the North of 1F Unit 5-6 Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at 1F South Discharge Channel (Bq/L)

