Reference

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station >

(Data summarized on November 29)

| Place of Sampling | North of Unit 5-6 Discharge Channel at Fukushima Daiichi NPS (Approx. 30m North of Unit 5-6 Discharge Channel) | | Around South Discharge Channel of Fukushima Daiichi NPS (Appox. 1.3km South of Unit 1-4 Discharge Channel) | | Specified by the Reactor Regulation (Bq/L) (The density limit in the water outside the surrounding monitored |
|----------------------------------|--|-------------------------|--|-------------------------|--|
| Time of Sampling | Nov 28, 2012 7:45 AM | | Nov 28, 2012 8:15 AM | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | areas is provided in section 6 of Appendix 2.) |
| I-131 (Approx. 8 days) | ND | - | ND | - | 40 |
| Cs-134 (Approx. 2 years) | ND | - | ND | - | 60 |
| Cs-137 (Approx. 30 years) | ND | - | ND | - | 90 |

* The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

* Data of other nuclides is under evaluation.


* In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.


* "ND" indicates that the measurement result is below the detection limit.

I-131: Approx. 0.50Bq/L, Cs-134: Approx.1.1Bq/L, Cs-137: Approx.1.5Bq/L

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detect

Radioactivity Density of the Seawater at 1F Units 5-6 North Discharge Channel (Bq/L)





Radioactivity Density of the Seawater at 1F Units 5-6 South Discharge Channel (Bq/L)

Sampling was conducted at around South Discharge Channel of Fukushima Daiichi NPS (appox. 330m south of Units 1-4 Discharge Channel) until November 25, 2012.