

## Result of Pu nuclide analysis in the soil Fukushima Daiichi Nuclear Power Station

### 1. Result analysis

( Unit : Bq/kg·dry soil )

Place of sampling ( ) shows distance from stuck of Unit 1/2	Date Analysis institute	Pu-238	Pu-239,Pu-240
Ground (WNW approx. 500m)	January 16 Japan Chemical Analysis Center	$( 1.5 \pm 0.13 ) \times 10^{-1}$	$( 5.9 \pm 0.77 ) \times 10^{-2}$
Yachounomori (W approx. 500m)		N.D. [ $<9.5 \times 10^{-3}$ ]	N.D. [ $<9.9 \times 10^{-3}$ ]
Around industrial waste treatment facility (SSW approx. 500m)		$( 2.9 \pm 0.20 ) \times 10^{-1}$	$( 1.1 \pm 0.11 ) \times 10^{-1}$
Domestic soil		N.D. $\sim 1.5 \times 10^{-1}$	N.D. $\sim 4.5$

[ ] shows lower detection limit

: Source: Ministry of Education, Culture, Sports, Science and  
Technology "Environmental radiation data base" from 1978 to 2008

: Place of sampling for " Ground" and " Around industrial waste treatment facility" has slightly  
changed to avoid duplication with past sampling and as for " Yachounomori", it was taken at  
the same point in depth direction (sampling point will be changed if sampling was not  
feasible).

### 2. Evaluation

Radioactive density of the Pu-239 and Pu-240 detected on January 16 was within the same  
level as that of fallout of past nuclear test in the atmosphere. However it is considerable the  
result may be derived from the nuclear accident this time.

Though there are some samples where Pu-238, Pu-239 and Pu-240 were detected after  
March 21, there is no significant change in the figures.

End