Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station >

(Data summarized on May 16)

Place of Sampling	North of Unit 5-6 Discharge Daiichi N (Approx. 30m North of Unit 8	IPS	Around South Discharge C Daiichi N (Appox. 1.3km South of Unit	the Reactor Regulation (Bq/L)			
Time of Sampling	May 15, 2 6:50 A		May 15, 2 7:05 A	(The density limit in the water outside the surrounding monitored areas is provided in			
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	section 6 of Appendix 2.)		
I-131 (Approx. 8 days)	ND	-	ND	-	40		
Cs-134 (Approx. 2 years)	ND	-	ND	-	60		
Cs-137 (Approx. 30 years)	ND	-	ND	ND -			

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

I-131: Approx. 0.42Bq/L, Cs-134: Approx. 1.1Bq/L, Cs-137: Approx. 1.5Bq/L

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

 $[\]ensuremath{^{*}}$ "ND" indicates that the measurement result is below the detection limit.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station, Remeasurement >

(Data summarized on May 16)

Place of Sampling	North of Unit 5-6 Discharge Daiichi N (Approx. 30m North of Unit 5	IPS	Around South Discharge C Daiichi N (Appox. 1.3km South of Unit	the Reactor Regulation (Bq/L)		
Time of Sampling	Apr 8, 20 6:55 A		Apr 8, 2 7:20 A	(The density limit in the water outside the surrounding monitored areas is provided in		
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	section 6 of Appendix 2.)	
Cs-134 (Approx. 2 years)	0.47	0.01	0.39	0.01	60	
Cs-137 (Approx. 30 years)	0.87	0.01	0.75	0.01	90	

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by: Tokyo Electric Power Environmental Engineering Co., Inc.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daiichi Nuclear Power Station, Remeasurement >

(Data summarized on May 16)

Place of Sampling	North of Unit 5-6 Discharge Daiichi N (Approx. 30m North of Unit 5	IPS	Around South Discharge C Daiichi N (Appox. 1.3km South of Unit	the Reactor Regulation (Bq/L)		
Time of Sampling	Apr 15, 2 6:40 A		Apr 15, 2 7:05 A	(The density limit in the water outside the surrounding monitored areas is provided in		
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	section 6 of Appendix 2.)	
Cs-134 (Approx. 2 years)	0.78 0.01		0.33	0.01	60	
Cs-137 (Approx. 30 years)	1.5	0.02	0.58	0.01	90	

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by: Tokyo Electric Power Environmental Engineering Co., Inc.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daini Nuclear Power Station >

(Data summarized on May 16)

Place of Sampling	2F Around the North D (Around Unit 3-4 Disc (Approx. 10km	charge Channel)	Around the North Sid (Approx. 11km South of I Chann (Approx. 23km	② Density Limit Specified by the Reactor Regulation (Bq/L)			
Time of Sampling	Apr 9, 2 9:30 A		Apr 9, 2 7:30 A	(The density limit in the water outside the surrounding monitored areas is provided in			
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	section 6 of Appendix 2.)		
I-131 (Approx. 8 days)	ND	-	ND	-	40		
Cs-134 (Approx. 2 years)	0.27	0.00	0.086	0.00	60		
Cs-137 (Approx. 30 years)	0.47	0.01	0.17	0.00	90		

^{*} The density specified by the Reactor Regulation is converted from Bg/cm³ to Bg/L.

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit. I-131: Approx. 0.50Bq/L

^{*} As to Cs-134 and Cs-137, analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted. Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Nuclides Analysis Result of the Radioactive Materials in the Seawater < Coast, Fukushima Daini Nuclear Power Station >

(Data summarized on May 16)

Place of Sampling	2F Around the North D (Around Unit 3-4 Disc (Approx. 10km	charge Channel)	Around the North Sid (Approx. 11km South of I Chann (Approx. 23km	② Density Limit Specified by the Reactor Regulation (Bq/L)		
Time of Sampling	Apr 16, 2 10:00 A		Apr 16, 2 7:20 A	(The density limit in the water outside the surrounding monitored areas is provided in		
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	section 6 of Appendix 2.)	
I-131 (Approx. 8 days)	ND	-	ND	-	40	
Cs-134 (Approx. 2 years)	0.11 0.00		0.058	0.00	60	
Cs-137 (Approx. 30 years)	0.20 0.00		0.11	0.00	90	

^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit. I-131: Approx. 0.46Bq/L

^{*} As to Cs-134 and Cs-137, analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted. Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Nuclides Analysis Result of Radioactive Materials in the Seawater < Offshore >

(Data summarized on May 16)

Place of Sampling (Place No.)	Зкт Оп	3km Offshore of Ukedo River (T-D1) Upper Layer Lower Layer			3km Offshore of Ukedo River (T-D1) Upper Layer Lower Layer				3km Offshore of Fukushima Daiichi NPS (T-D5) Upper Layer Lower Layer				② Density Limit Specified by the Reactor Regulation (Bg/L)	
Time of Sampling	Anr 9 20	013	Apr 9, 20 9:04 Al	013	Apr 16, 2 8:58 A	2013	Apr 16, 2 8:58 Al	013	Apr 9, 20	Apr 9, 2013 Apr 9, 2 9:33 AM 9:33 A		013	(The density limit in the	
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	araga ia providad in	
Cs-134 (Approx. 2 years)	0.032	0.00	0.031	0.00	0.025	0.00	0.022	0.00	0.034	0.00	0.021	0.00	60	
Cs-137 (Approx. 30 years)	0.062	0.00	0.055	0.00	0.052	0.00	0.044	0.00	0.062	0.00	0.040	0.00	90	

Place of Sampling (Place No.)	3km Offshore of Fukushima Daiichi NPS (T-D5) Upper Layer Lower Layer				3km Offshore of Fukushima Daini NPS (T-D9) Upper Layer Lower Layer				3km Offshore of Fukushima Daini NPS (T-D9)				② Density Limit Specified by the Reactor Regulation
Time of Sampling	Anr 16 2	013	Apr 16, 2 8:26 Al	013	Apr 10, 2 8:52 Al	013	Apr 10, 2 8:52 Al	013	Upper La Apr 17, 2 8:53 Al	013 Apr 17, 2013		(Bq/L) (The density limit in the water outside the surrounding monitored	
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	areas is provided in section 6 of Appendix 2.)
Cs-134 (Approx. 2 years)	0.037	0.00	0.037	0.00	0.022	0.00	0.024	0.00	0.013	0.00	0.063	0.00	60
Cs-137 (Approx. 30 years)	0.078	0.00	0.074	0.00	0.047	0.00	0.042	0.00	0.026	0.00	0.12	0.00	90


^{*} The density specified by the Reactor Regulation is converted from Bq/cm³ to Bq/L.

 $^{^{\}star}$ In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.


^{*} Analysis results by detail analysis (Phosphomolybdic acid ammonium adsorption sampling method) are noted.

^{*} Analyzed by: Tokyo Electric Power Environmental Engineering Co., Inc.


Radioactivity Density of the Seawater at 1F Units 5-6 North Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at 1F South Discharge Channel (Bq/L)



Radioactivity Density of the Seawater at 2F North Discharge Channel (Bq/L)



Radioactivity Density of the Seawater Around the South Side of Kitasakogawa (Bq/L)

