Reference

Nuclide Analysis Results of the Radioactive Materials in the Air at Fukushima Nuclear Power Stations < 1/2 >

(Data summarized on January 15)

Place of Sampling	The West Gate of Daiichi N						② Density Limit Specified by the Reactor Regulation
Time of Sampling	January 14, 2014 7:00 AM - 12:00 PM						(Bq/cm³) (Density limit in the air which radiation workers breathe in
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	is specified in section 4 of Appendix 2)
I-131 (Approx. 8 days)	ND	-					1E-03
Cs-134 (Approx. 2 years)	ND	-					2E-03
Cs-137 (Approx. 30 years)	ND	-					3E-03

^{*} The radioactivity density is the sum of the volatile nuclides density and the particulate nuclides density.

O.OE-O is the same as O.O x 10 $^{-}$ O

Data of other nuclides is under examination.

The detection limits at the west gate of Fukushima Daiichi NPS are as follows: Volatile: I-131: Approx. 7E-8Bq/cm3, Cs-134: Approx.2E-7Bq/cm3, Cs-137: Approx.2E-7Bq/cm3 Particulate: I-131: Approx. 5E-8Bq/cm3, Cs-134: Approx.9E-8Bq/cm3, Cs-137: Approx.1E-7Bq/cm3 As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.

Reference

Nuclide Analysis Results of the Radioactive Materials in the Air at Fukushima Nuclear Power Stations < 2/2 >

(Data summarized on January 15)

Place of Sampling	MP-1 at Fukushima Daiichi NPS		MP-3 at Fukushima Daiichi NPS		MP-8 at Fukushima Daiichi NPS		② Density Limit Specified by the Reactor Regulation (Bq/cm³) (Density limit in the air which radiation workers breathe in
Time of Sampling	January 14, 2014 8:14 AM - 1:14 PM		January 14, 2014 8:48 AM - 1:48 PM		January 14, 2014 8:26 AM - 1:26 PM		
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	①Density of Sample (Bq/cm³)	Scaling Factor (①/②)	is specified in section 4 of Appendix 2)
I-131 (Approx. 8 days)	ND	-	ND	-	ND	-	1E-03
Cs-134 (Approx. 2 years)	ND	-	ND	-	ND	-	2E-03
Cs-137 (Approx. 30 years)	ND	-	ND	-	ND	-	3E-03

^{*} The radioactivity density is the sum of the volatile nuclides density and the particulate nuclides density.


O.OE-O is the same as $O.O \times 10^{-O}$


Data of other nuclides is under examination.


The detection limits are as follows. Volatile: I-131: Approx. 6E-8Bq/cm3, Cs-134: Approx.7E-8Bq/cm3, Cs-137: Approx.7E-8Bq/cm3
Particulate: I-131: Approx. 4E-8Bq/cm3, Cs-134: Approx.4E-8Bq/cm3, Cs-137: Approx.4E-8Bq/cm3
As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

^{*} In the case of more than 2 nuclides, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.







Dust Nuclides Analysis Result: MP-8 at Fukushima Daiichi NPS (Bq/cm³)

