Underground Reservoir Nuclide Analysis Results (As of June 23, 2014)

| | | | | | | U | ndergrour | nd Reserv | oir (Drain | hole wate | er) | | | | |
|-----------------------|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | | i | | ii | | iii | | iv | | ٧ | | vi | | \ | vii |
| | | Northeast side | Southwest side |
| Sam | Sampled time | | / | 7:52 AM | / | 7:42 AM | 7:45 AM | / | / | / | / | / | | / | |
| Chloride cor | chloride concentration (ppm) | | | 8 | | 5 | 3 | | | | | | | | |
| | I-131 | <2.7E-2 | | <1.9E-2 | | <2.1E-2 | <2.2E-2 | | | | | | | | |
| Radioactive | Cs-134 | <4.7E-2 | | <3.8E-2 | | <4.3E-2 | <3.9E-2 | | | | | | | | |
| concentration | Cs-137 | <6.4E-2 | | <6.1E-2 | | <6.4E-2 | <5.8E-2 | | | | | | | | |
| | γ nuclides other than the major 3 nuclides | ND | | ND | | ND | ND | | | | | | | | |
| (Bq/cm ³) | ΑΙΙ β | 2.0E-1 | / | 4.3E-2 | / | 7.6E-2 | <3.0E-2 | / | / | / | / | / | / | / | / |

Half-life period I-131: Approx. 8 days, Cs-134: Approx. 2 years, Cs-137: Approx. 30 years

| | | Underground Reservoir (Leakage detector hole water) | | | | | | | | | | | | | |
|------------------------------|--|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | | i | | ii | | iii | | iv | | v / | | vi | | vii / | |
| | | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side | Northeast side | Southwest side |
| Sampled time | | 7:30 AM | / | 7:26 AM | / | 7:37 AM | 7:48 AM | / | | | | / | | | |
| Chloride concentration (ppm) | | 10 | | 15 | | 9 | 9 | | | | | | | | |
| | I-131 | <2.6E-2 | | <2.2E-2 | | <2.2E-2 | <2.3E-2 | | | / | ľ | | | / | |
| Radioactive | Cs-134 | <4.3E-2 | | <4.3E-2 | | <4.3E-2 | <4.1E-2 | | | | | | | | |
| concentration | Cs-137 | <6.4E-2 | | <6.0E-2 | | <6.4E-2 | <5.6E-2 | | | | | | | | |
| | γ nuclides other than the major 3 nuclides | ND | | ND | | ND | ND | | | | | | | | |
| (Bq/cm ³) | ΑΙΙ β | 5.9E+1 | | 1.7E+1 | | 2.1E+1 | 9.6E+0 | / | | | | | / | | |

Half-life period I-131: Approx. 8 days, Cs-134: Approx. 2 years, Cs-137: Approx. 30 years

(Note 1) O.OE \pm O is the same as O.O x $10^{\pm O}$.

(Note 2) The figures written next to "<" indicate the detection limit when the measurement result is below the detection limit.

(Note 3) "ND" indicates that the measurement result of γ nuclides other than the major 3 nuclides are below the detection limit.

Nuclide Analysis Results of the Underground Bypass (Investigation Holes/Pumping Well) and the Sea Side Observation Holes (As of June 23, 2014)

| | | erground by estigation he | - | Sea side observation holes | | | | | | | | |
|-------------------------------|---|------------------------------|---|----------------------------|---|---|---|-------------------|-------------------|-------------------|-------------------|--|
| | а | b | O | 1 | 2 | 3 | 4 | (5) | 6 | 7 | 8 | |
| Sampled time | | | | | | | | 9:43 AM | 9:26 AM | 10:03 AM | 9:08 AM | |
| Chloride concentration (ppm) | | | | | | | | 8 | 9 | 13 | 11 | |
| Tritium (Bq/cm ³) | | | | | | | | Under analysis | Under analysis | Under analysis | Under analysis | |
| All β(Bq/cm ³) | | | | | | | | <3.0E-2 | <3.0E-2 | <3.0E-2 | <3.0E-2 | |

Half-life period Tritium: Approx. 12 years

(Note 1) O.OE \pm O is the same as O.O x $10^{\pm O}$.

(Note 2) The figures written next to "<" indicate the detection limit when the measurement result is below the detection limit.