Reference

Radioactivity Density of the Seawater in the Port of Fukushima Daiichi NPS<1/2>

(Data summarized on December 4)

Place of Sampling	Sha	allow Draft	Quay at 1F *		Inside Unit 1-4 Water Intake Canal (North) at Fukushima Daiichi NPS (North side of the East Seawall Break)				Inside Unit 1-4 Water Intake Canal (South) at Fukushima Daiichi NPS (in front of Impermeable Wall)				② Density Limit Specified by the Reactor Regulation
Time of Sampling	Dec 3, 2014 7:02 AM		N/A		Dec 3, 2014 7:30 AM		Dec 3, 2014 7:22 AM		Dec 3, 2014 7:25 AM		N/A		(Bq/L) (The density limit in the water outside the
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (1/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	surrounding monitored areas is provided in section 6 of Appendix 2.)
I-131 (Approx. 8 days)	ND	-	-	-	ND	-	ND	-	ND	-	-	-	40
Cs-134 (Approx. 2 years)	ND	-	-	-	3.2	0.05	11	0.18	4.1	0.07	-	-	60
Cs-137 (Approx. 30 years)	ND	-	-	-	11	0.12	38	0.42	18	0.20	-	-	90

^{*} The density specified by the Reactor Regulation is converted from Bq/cm^3 to Bq/L.

^{*} Data of other nuclides is under evaluation.

^{*} In the case of 2 nuclides or more, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.

I-131: Approx. 3Bq/L, Cs-134: Approx.2Bq/L, Cs-137: Approx.2Bq/L As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected. * The sampling will be performed after opening and closing of the silt fence.

Reference

Radioactivity Density of the Seawater in the Port of Fukushima Daiichi NPS<2/2>

(Data summarized on December 4)

Place of Sampling	Port Entrar	ice of Fuki	ushima Daiichi NI	PS*									② Density Limit Specified by the Reactor Regulation
Time of Sampling	Dec 2, 2014 1:40 PM		N/A										(Bq/L) (The density limit in the water outside the
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	surrounding monitored areas is provided in section 6 of Appendix 2.)										
I-131 (Approx. 8 days)	ND	-	-	-									40
Cs-134 (Approx. 2 years)	ND	-	-	-									60
Cs-137 (Approx. 30 years)	1.3	0.01	-	-									90

^{*} The density specified by the Reactor Regulation is converted from Bq/cm^3 to Bq/L. * Data of other nuclides is under evaluation.


^{*} In the case of 2 nuclides or more, the sum of scaling factors to density limits is compared to 1.

^{* &}quot;ND" indicates that the measurement result is below the detection limit.


As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected. * I-131: Approx. 1Bg/L, Cs-134: Approx.1Bg/L

^{*} At these points, sampling is carried out once a week. (As for the port entrance, also sampled on the day the silt fence was opened/shut or covering work was carried out in the port.)


Radioactivity Density of the Seawater in Front of the Shallow Draft Quay at 1F (Bq/L)




Radioactivity Density of the Seawater at the North of Unit 1-4 Water Intake (North of East Seawater Break of Fukushima Daiichi NPS (Bq/ L)



Radioactivity Density of the Seawater at Unit 4 Screen at Fukushima Daiichi NPS (Bq/L)



Radioactivity Density of the Seawater at the South of Unit 1-4 Water Intake (in front of Impermeable Wall) at Fukushima Daiichi NPS (Bq/L)



Radioactivity Density of the Seawater at the Port Entrance of Fukushima Daiichi NPS (Bq/L)

