Analyzed result of nuclide in fish < Whitin the port at Fukushima Daiichi NPŚ>

(Data summarized on September 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish(muscle)	Whitin the port (Near south breakwater) in Fukushima Daiichi NPS	Aug 5, 2015	13	80	93
Greenling(muscle) No.1	Whitin the port (Near port entrance) in Fukushima Daiichi NPS	Aug 1, 2015	38	120	158
Spotted halibut(muscle)	Whitin the port (Near port entrance) in Fukushima Daiichi NPS	Aug 1, 2015	59	250	309
Greenling(muscle) No.2	Whitin the port (Near port entrance) in Fukushima Daiichi NPS	Aug 3, 2015	140	660	800
Schlegel's black rockfish(muscle)	Whitin the port (Near port entrance) in Fukushima Daiichi NPS	Aug 3, 2015	75	290	365
Greenling(muscle) No.3	Whitin the port (Near port entrance) in Fukushima Daiichi NPS	Aug 5, 2015	290	1100	1390
Marbled sole(muscle)	Whitin the port (Near port entrance) in Fukushima Daiichi NPS	Aug 5, 2015	260	1000	1260

^{*} When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

 $^{^{\}star}\,$ Standard value (Since April 1, 2012) The total of Cs-134 and Cs-137: 100Bq/kg

^{*} Analyzed by: Tokyo Power Technology Ltd.