

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) <1/6>
 (excluding inside the port)

(Data summarized on March 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137(Approx. 30 years)	CS (Total)
Sebastes cheni(muscle)	Around 1km Offshore of Ota River (T-S1)	February 27, 2015	7.5	27	34.5
Schlegel's black rockfish(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	ND(4.1)	15	15
Common skete(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	6.2	18	24.2
Sebastes cheni(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	16	50	66
Sea bass(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	ND(3.7)	ND(3.7)	ND
Microstomus achne(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	13	56	69
Ovalipes punctatus(whole)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	ND(3.7)	ND(4.0)	ND
Flatfish(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	ND(3.4)	ND(4.3)	ND
Marbled sole(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	ND(3.5)	6.7	6.7
Pacific cod(muscle)	Around 3km Offshore of Odaka Ward (T-S2)	February 27, 2015	ND(3.3)	ND(4.5)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) <2/6>
(excluding inside the port)

(Data summarized on March 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137(Approx. 30 years)	CS (Total)
Northern dogfish(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(3.7)	ND(4.2)	ND
Schlegel's black rockfish(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(3.7)	ND(4.2)	ND
Sea raven(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(4.6)	5.2	5.2
Common skete(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	10	28	38
Ovalipes punctatus(whole)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(3.6)	4.1	4.1
Flatfish(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(4.0)	5.7	5.7
Marbled sole(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(4.5)	12	12
Pacific cod(muscle)	Around 3km Offshore of Ukedo River (T-S3)	February 25, 2015	ND(3.4)	3.9	3.9
Northern dogfish(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(3.0)	ND(3.8)	ND
Blue crab(whole)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(3.2)	ND(3.5)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) <3/6>
(excluding inside the port)

(Data summarized on March 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137(Approx. 30 years)	CS (Total)
Sea raven(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(3.8)	ND(4.3)	ND
Common skete(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(4.2)	17	17
Microstomus achne(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	6.3	24	30.3
Ovalipes punctatus(whole)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(3.5)	ND(3.7)	ND
Flatfish(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(4.1)	10	10
Marbled sole(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	5.6	18	23.6
Pacific cod(muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	February 25, 2015	ND(4.3)	4.8	4.8
Sea raven(muscle)	Around 2km Offshore of Kido River (T-S5)	February 13, 2015	ND(4.2)	ND(4.0)	ND
Common skete(muscle)	Around 2km Offshore of Kido River (T-S5)	February 13, 2015	12	41	53
Flatfish(muscle)	Around 2km Offshore of Kido River (T-S5)	February 13, 2015	4.3	11	15.3

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 4/6 >
(excluding inside the port)

(Data summarized on March 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137(Approx. 30 years)	CS (Total)
Marbled sole(muscle)	Around 2km Offshore of Kido River (T-S5)	February 13, 2015	6.9	20	26.9
Pacific cod(muscle)	Around 2km Offshore of Kido River (T-S5)	February 13, 2015	ND(3.9)	ND(4.5)	ND
Sea raven(muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	February 13, 2015	ND(4.3)	14	14
Common skete(muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	February 13, 2015	9.9	33	42.9
Flatfish(muscle)	Offshore 2km vicinity of Fukushima Daiichi NPS (T-S7)	February 13, 2015	16	52	68
Ovalipes punctatus(whole)	Around 4km Offshore of Kumagawa (T-S8)	February 09, 2015	ND(3.7)	ND(4.0)	ND
Flatfish(muscle)	Around 4km Offshore of Kumagawa (T-S8)	February 09, 2015	ND(3.1)	ND(3.6)	ND
Stone flounder(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(3.8)	ND(3.4)	ND
Common skete(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(3.3)	7.1	7.1
Sea bass(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(3.6)	ND(3.9)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) <5/6>
(excluding inside the port)

(Data summarized on March 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137(Approx. 30 years)	CS (Total)
Microstomus achne(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(3.5)	ND(4.3)	ND
Flatfish(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(4.0)	ND(3.7)	ND
Littlemouth flounder(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(3.2)	5.3	5.3
Roundnose flounder(muscle)	Around 15km Offshore of Odaka Ward (T-B1)	February 25, 2015	ND(4.1)	ND(3.7)	ND
Greenling(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.3)	ND(4.0)	ND
Stone flounder(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.6)	4.3	4.3
Common skete(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	8.4	27	35.4
Sea bass(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.5)	ND(3.6)	ND
Microstomus achne(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.5)	ND(3.8)	ND
Flatfish(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(4.1)	ND(3.4)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) <6/6>
(excluding inside the port)

(Data summarized on March 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137(Approx. 30 years)	CS (Total)
Littlemouth flounder(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.4)	ND(4.5)	ND
Marbled sole(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(4.2)	5.2	5.2
Pacific cod(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.6)	ND(3.7)	ND
Roundnose flounder(muscle)	Around 18km Offshore of Ukedo River (T-B2)	February 25, 2015	ND(3.7)	ND(3.9)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by: Tokyo Power Technology Ltd.