

Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on September 21)

| Name of Sample (Region) | Place of Sampling (Place No.) | Date of Sampling | Radioactivity Concentration [Bq/kg (Raw)] (Half-life) | | |
|---------------------------------|--|------------------|---|------------------------------|----------|
| | | | Cs-134 (Approx. 2 years) | Cs-137 (Approx. 30 years) | CS (Sum) |
| Acanthopagrus schlegeli(muscle) | Port area of Fukushima Daiichi NPS (Near the port entrance) | Aug.3, 2016 | ND(7.8) | 19 | 19 |
| Flatfish(muscle) | Port area of Fukushima Daiichi NPS (Near the port entrance) | Aug.3, 2016 | ND(8.7) | 51 | 51 |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.