

Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on October 18)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Acanthopagrus schlegeli (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 2, 2016	12	91	103
Common skate (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 2, 2016	13	96	109
Greenling (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 7, 2016	27	160	187
Drumfish (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 7, 2016	ND (8.1)	12	12
Schlegel's black rockfish (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 27, 2016	8.9	49	57.9
Seriola dumerili (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 28, 2016	ND (5.3)	32	32
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 28, 2016	180	1100	1280
Chum salmon (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 29, 2016	ND (5.5)	7.7	7.7
Flathead (Platycephalus sp.) (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Sept. 29, 2016	ND (5.6)	9.4	9.4

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.