Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on November 22)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle) No.1	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 7, 2016	23	190	213
Striped mullet (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 7, 2016	ND(7.2)	23	23
Common skete (muscle) No.1	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 13, 2016	ND(7.0)	27	27
Sebastes cheni (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 13, 2016	650	3900	4550
Flatfish (muscle) No.2	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 13, 2016	7.5	63	70.5
Yellowtail (muscle)	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 13, 2016	ND(7.1)	ND(7.2)	ND
Chum salmon (muscle) No.1	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 19, 2016	ND(5.4)	ND(5.5)	ND
Common skete(muscle) No.2	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 28, 2016	8.7	50	58.7
Chum salmon (muscle) No.2	Port area of Fukushima Daiichi NPS (Near the port entrance)	Oct. 28, 2016	ND(6.5)	ND(6.2)	ND

^{*}When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

^{*}Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.