

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar 24, 2016	ND (4.0)	6.1	6.1
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar 24, 2016	ND (4.0)	ND (3.8)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar 24, 2016	ND (4.1)	ND (4.1)	ND
Pacific cod (muscle)	Around 1km Offshore of Ota River (T-S1)	Mar 24, 2016	ND (4.2)	ND (4.1)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar 24, 2016	ND (3.5)	11	11
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar 24, 2016	ND (4.0)	ND (4.4)	ND
Pacific cod (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Mar 24, 2016	ND (3.9)	ND (3.9)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar 25, 2016	ND (2.9)	4.8	4.8
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar 25, 2016	ND (3.5)	5.3	5.3
Microstomus achne (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar 25, 2016	ND (4.4)	13	13

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar 25, 2016	ND (3.6)	5.8	5.8
Pacific cod (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Mar 25, 2016	ND (3.7)	ND (4.2)	ND
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Mar 25, 2016	ND (3.2)	ND (3.5)	ND
Common skate (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Mar 25, 2016	ND (4.5)	21	21
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Mar 25, 2016	ND (3.5)	15	15
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Mar 25, 2016	ND (3.6)	7.9	7.9
Pacific cod (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Mar 25, 2016	ND (4.1)	ND (3.8)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	Mar 25, 2016	ND (3.2)	ND (3.6)	ND
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (4.5)	7.1	7.1
Northern dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (3.7)	ND (3.4)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea raven (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (3.5)	5.9	5.9
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (4.4)	20	20
Sebastes cheni (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	10	45	55
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	6.6	27	33.6
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (4.1)	ND (3.6)	ND
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (3.1)	15	15
Pacific cod (muscle)	Around 2km Offshore of Kido River (T-S5)	Mar 17, 2016	ND (3.6)	ND (3.8)	ND
Greenling (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Mar 17, 2016	ND (3.4)	6.6	6.6
Common skete (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Mar 17, 2016	ND (3.3)	10	10
Microstomus achne (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Mar 17, 2016	ND (3.1)	9.0	9.0

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Mar 17, 2016	ND (4.1)	6.5	6.5
Pacific cod (muscle)	Offshore 2km vicinity of Fukushima Daini NPS (T-S7)	Mar 17, 2016	ND (3.6)	ND (3.9)	ND
Stone flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (3.5)	13	13
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (3.9)	ND (3.7)	ND
Schlegel's black rockfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (4.2)	ND (3.7)	ND
Common skate (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (4.1)	8.2	8.2
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (3.5)	9.7	9.7
Ovalipes punctatus (whole)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (3.2)	ND (3.7)	ND
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (4.2)	6.0	6.0
Pacific cod (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Mar 7, 2016	ND (4.2)	ND (3.7)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.5)	ND (4.5)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (2.7)	ND (3.8)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.2)	ND (3.6)	ND
Common skate (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.6)	6.2	6.2
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.9)	ND (3.8)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (4.3)	4.3	4.3
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (4.0)	ND (4.0)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.5)	ND (3.7)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	7.0	34	41
Pacific cod (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.7)	ND (3.7)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (4.2)	ND (3.2)	ND
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Mar 3, 2016	ND (3.1)	ND (3.8)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (2.9)	12	12
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.8)	ND (3.6)	ND
Common skate (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (4.5)	ND (4.1)	ND
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.8)	ND (3.8)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.1)	ND (3.7)	ND
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (4.5)	ND (3.3)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.4)	ND (4.0)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.6)	7.2	7.2

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.5)	ND (3.2)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (3.4)	ND (3.2)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Mar 3, 2016	ND (4.2)	ND (4.0)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (3.6)	ND (4.4)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (3.7)	16	16
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (3.3)	ND (3.9)	ND
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (4.3)	11	11
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (4.0)	ND (4.4)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (4.1)	8.3	8.3
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (4.1)	ND (4.1)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (2.9)	ND (4.1)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (3.0)	ND (3.5)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (4.2)	7.4	7.4
Pacific cod (muscle)	Around 10km Offshore of Fukushima Daiichi NPS (T-B3)	Mar 25, 2016	ND (4.1)	ND (4.0)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (3.9)	ND (3.9)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (3.6)	9.7	9.7
Sea bass (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (3.2)	ND (4.1)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (3.6)	6.3	6.3
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (3.6)	ND (3.5)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (4.0)	4.2	4.2

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish (Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/9>
 (excluding the Port Area)

(Data summarized on April 15)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (4.0)	ND (4.3)	ND
Pacific cod (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (2.9)	ND (3.6)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	Mar 25, 2016	ND (3.7)	ND (3.7)	ND

* When analyzed result is less than a detection limit level of the radioactivity concentration, showed as "ND"; as of individual Nuclide, detection limit level is showed in parenthesis.

* The sum of Cs-134 and Cs-137 radioactivity concentration as a standard value (since April 1, 2012) is 100 Bq/kg.

* Analyzed by Tokyo Power Technology Ltd.