

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(3.7)	ND(4.0)	ND
Acanthopagrus schlegeli (muscle)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(3.7)	12	12
Common skate (muscle)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(3.1)	4.4	4.4
Chum salmon (muscle)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(4.2)	ND(4.0)	ND
Drumfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(3.6)	ND(4.0)	ND
Flatfish① (muscle)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(3.6)	ND(3.7)	ND
Flatfish② (muscle)	Around 1km Offshore of Ota River (T-S1)	Nov. 11, 2016	ND(3.4)	ND(3.6)	ND
Stingray (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(4.1)	ND(4.0)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.4)	ND(3.9)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.3)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Angel shark (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.6)	16	16
Acanthopagrus schlegeli (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.9)	ND(3.2)	ND
Sea raven (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(4.3)	ND(3.9)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.6)	ND(3.3)	ND
Chum salmon (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.6)	ND(3.8)	ND
Sebastes cheni (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(4.1)	14	14
Drumfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(4.2)	3.5	3.5
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.8)	3.9	3.9
Flatfish① (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.9)	4.3	4.3
Flatfish② (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.7)	ND(2.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(4.0)	ND(3.5)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.9)	ND(3.2)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.0)	6.8	6.8
Pagrus major (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Nov. 11, 2016	ND(3.5)	ND(3.9)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(4.2)	ND(3.7)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(3.7)	ND(4.1)	ND
Common skate (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(3.3)	14	14
Drumfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(3.0)	ND(3.5)	ND
Microstomus achne (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(3.7)	9.1	9.1
Flatfish① (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(3.7)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish② (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(4.2)	ND(4.0)	ND
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Nov. 11, 2016	ND(3.8)	7.5	7.5
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Nov. 11, 2016	ND(3.6)	ND(3.8)	ND
Blue crab (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Nov. 11, 2016	ND(4.1)	ND(3.8)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Nov. 11, 2016	ND(4.1)	11	11
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Nov. 11, 2016	ND(3.9)	ND(3.7)	ND
Sea robin (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Nov. 11, 2016	ND(3.6)	ND(3.8)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Nov. 11, 2016	ND(3.8)	8.9	8.9
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(3.3)	ND(4.2)	ND
Angel shark (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(4.4)	3.8	3.8

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Acanthopagrus schlegeli (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(3.8)	ND(3.5)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(3.3)	9.3	9.3
Drumfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(3.3)	3.5	3.5
Flatfish① (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(3.4)	ND(3.6)	ND
Flatfish② (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(4.0)	5.1	5.1
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(3.4)	4.7	4.7
Flathead (Platycephalus sp.) (muscle)	Around 2km Offshore of Kido River (T-S5)	Nov. 14, 2016	ND(4.0)	ND(3.7)	ND
Stingray (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Nov. 14, 2016	ND(3.8)	ND(3.1)	ND
Angel shark (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Nov. 14, 2016	ND(4.1)	5.8	5.8
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Nov. 14, 2016	ND(3.8)	11	11

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Banded dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Nov. 14, 2016	ND(3.6)	14	14
Flatfish① (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Nov. 14, 2016	ND(3.9)	4.2	4.2
Flatfish② (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Nov. 14, 2016	ND(4.4)	4.3	4.3
Stingray (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(3.3)	4.1	4.1
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(3.9)	ND(2.8)	ND
Angel shark (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	6.8	45	51.8
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(3.4)	8.4	8.4
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(3.7)	16	16
Flatfish① (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(3.8)	4.0	4.0
Flatfish② (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(4.0)	4.1	4.1

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Nov. 16, 2016	ND(3.7)	ND(3.6)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(3.1)	ND(2.9)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(3.3)	ND(3.6)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(4.2)	ND(4.0)	ND
Yellowtail (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(3.5)	ND(3.9)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(3.7)	ND(4.2)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(3.9)	ND(4.0)	ND
Pagrus major (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(4.0)	ND(3.9)	ND
Dory (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Nov. 18, 2016	ND(3.3)	4.8	4.8
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.7)	ND(3.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Striped jewfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(4.4)	ND(3.3)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.3)	ND(3.5)	ND
Lophius litilon (whole)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.6)	ND(3.1)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.6)	4.0	4.0
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.3)	ND(3.8)	ND
Cutlass fish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(4.4)	ND(3.6)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.7)	4.8	4.8
Flatfish① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.9)	ND(3.1)	ND
Flatfish② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.4)	ND(4.0)	ND
Smooth dogfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.0)	ND(3.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.3)	ND(3.6)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.2)	3.1	3.1
Dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.7)	ND(3.7)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.4)	ND(3.5)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Nov. 18, 2016	ND(3.6)	ND(3.4)	ND
Stingray (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(3.9)	ND(4.0)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(3.7)	ND(3.7)	ND
Angel shark (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(3.3)	ND(4.6)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(3.7)	ND(2.9)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(4.4)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Takifugu snyderi (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(3.4)	ND(3.2)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(4.0)	3.8	3.8
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(2.9)	ND(4.0)	ND
Pagrus major (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(2.8)	ND(3.5)	ND
Dory (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Nov. 7, 2016	ND(3.4)	ND(3.6)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.8)	4.3	4.3
Common skate (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.4)	5.4	5.4
Takifugu snyderi (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.9)	ND(3.7)	ND
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.4)	ND(3.6)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.6)	ND(3.8)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/11>
(excluding the port)

(Data summarized on December 20)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea robin (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.5)	ND(3.5)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.9)	ND(3.9)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(4.1)	ND(4.4)	ND
Pagrus major (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.8)	ND(4.1)	ND
Dory (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(3.6)	ND(3.9)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Nov. 7, 2016	ND(4.0)	ND(4.1)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.