Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on May 19)

				<u> </u>	
Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Schlegel's black rockfish (muscle)	Port area of Fukushima Daiichi NPS(Near the east seawall break)	Apr. 13, 2017	38	260	298
Greenling (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	Apr. 6, 2017	54	390	444
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	Apr. 6, 2017	ND(5.6)	39	39
Drumfish (muscle)	Port area of Fukushima Daiichi NPS(Near the port center)	Apr. 13, 2017	ND(4.4)	11	11

^{*}When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in paths as the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.