

## Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Drumfish (muscle) No.1	Port area of Fukushima Daiichi NPS(Near the east seawall break)	May 11, 2017	ND(5.8)	9.2	9.2
Drumfish (muscle) No.2	Port area of Fukushima Daiichi NPS(Near the north breakwater)	May 11, 2017	ND(4.3)	9.6	9.6
Greenling (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	May 12, 2017	ND(7.3)	62	62
Drumfish (muscle) No.3	Port area of Fukushima Daiichi NPS(Near the port entrance)	May 12, 2017	6.0	45	51
Drumfish (muscle) No.4	Port area of Fukushima Daiichi NPS(Near the port center)	May 25, 2017	ND(4.2)	11	11
/	/	/	/	/	/
/	/	/	/	/	/
/	/	/	/	/	/
/	/	/	/	/	/

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in p

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.