

Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

(Data summarized on August 17)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Drumfish (muscle) No.1	Port area of Fukushima Daiichi NPS(Near the east seawall break)	July 4, 2017	ND(6.1)	18	18
Konosirus punktatus (muscle)	Port area of Fukushima Daiichi NPS(Near the south breakwater)	July 20, 2017	ND(6.8)	58	58
Drumfish (muscle) No.2	Port area of Fukushima Daiichi NPS(Near the port entrance)	July 4, 2017	ND(4.7)	17	17
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS(Near the port entrance)	July 4, 2017	6.0	34	40
Drumfish (muscle) No.3	Port area of Fukushima Daiichi NPS(Near the port entrance)	July 10, 2017	ND(5.7)	13	13
/	/	/	/	/	/
/	/	/	/	/	/
/	/	/	/	/	/
/	/	/	/	/	/

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in p

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.