

Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Black sea bream (muscle) No.1	Port area of Fukushima Daiichi NPS(Near the east seawall break)	September 22, 2017	ND(4.5)	12	12
Black Sea bream (muscle) No.2	Port area of Fukushima Daiichi NPS(Near the east seawall break)	September 22, 2017	ND(7.4)	ND(7.9)	ND
Mugil cephalus (muscle) No.1	Port area of Fukushima Daiichi NPS(Near the east seawall break)	September 22, 2017	6.7	49	55.7
Greater amberjack (muscle) No.1	Port area of Fukushima Daiichi NPS(Near the north breakwater)	September 6, 2017	ND(5.4)	12	12
Greater amberjack (muscle) No.2	Port area of Fukushima Daiichi NPS(Near the north breakwater)	September 6, 2017	ND(4.7)	ND(5.2)	ND
Flatfish (muscle)	Port area of Fukushima Daiichi NPS(Near the north breakwater)	September 6, 2017	6.5	50	56.5
Konosirus punktatus (muscle)	Port area of Fukushima Daiichi NPS(Near the port center)	September 22, 2017	12	99	111
Mugil cephalus (muscle) No.2	Port area of Fukushima Daiichi NPS(Near the port center)	September 22, 2017	11	110	121

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in page 10.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.