

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stingray (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(3.6)	ND(3.9)	ND
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(4.4)	ND(4.1)	ND
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(3.9)	3.9	3.9
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(3.9)	5.4	5.4
Ovalipes punctatus (whole)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(3.6)	ND(4.5)	ND
Smooth dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(2.5)	ND(3.9)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	Jan. 26, 2017	ND(4.0)	6.9	6.9
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(4.0)	ND(4.4)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(4.0)	14	14
Sea raven (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(3.1)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(4.0)	4.7	4.7
Alaska pollack (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(4.2)	ND(3.6)	ND
Sea bass (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(3.2)	7.9	7.9
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(4.2)	6.9	6.9
Smooth dogfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(4.0)	ND(4.1)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	Jan. 26, 2017	ND(3.6)	4.4	4.4
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.2)	4.6	4.6
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(3.9)	ND(4.0)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.2)	ND(4.5)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(3.8)	ND(4.1)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.0)	19	19
Sea bass (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.5)	4.8	4.8
Ovalipes punctatus (whole)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(3.8)	ND(3.6)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(3.7)	ND(3.6)	ND
Dasyatis matsubara (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(3.5)	ND(3.5)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.5)	ND(3.1)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(3.7)	5.3	5.3
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.1)	4.7	4.7
Roundnose flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	Jan. 19, 2017	ND(4.0)	3.4	3.4
Blue crab (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(3.6)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Lophius litilon (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(4.3)	ND(3.7)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(4.1)	ND(4.6)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(3.6)	5.4	5.4
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(4.1)	3.8	3.8
Smooth dogfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(3.8)	ND(3.7)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(4.0)	ND(4.0)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(3.4)	ND(3.7)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	Jan. 19, 2017	ND(4.1)	ND(4.2)	ND
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(3.9)	6.5	6.5
Lophius litilon (whole)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(3.5)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Acanthopagrus schlegeli (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	7.2	43	50.2
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(3.4)	13	13
Banded dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(4.4)	7.3	7.3
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(4.1)	10	10
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(3.5)	ND(4.2)	ND
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(3.6)	5.5	5.5
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	Jan. 28, 2017	ND(3.4)	ND(3.7)	ND
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.9)	4.0	4.0
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(4.7)	ND(4.0)	ND
Lophius litilon (whole)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(4.0)	ND(4.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sebastes vulpes (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	4.3	24	28.3
Sea raven (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.4)	ND(4.3)	ND
Common skate (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.5)	11	11
Sebastes cheni (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	6.6	41	47.6
Banded dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.9)	5.0	5.0
Drumfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.8)	ND(3.6)	ND
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(4.3)	9.6	9.6
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.9)	ND(4.8)	ND
Smooth dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(3.6)	ND(3.8)	ND
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(4.0)	12	12

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	Jan. 28, 2017	ND(2.9)	ND(3.7)	ND
Stingray (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.9)	5.5	5.5
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.6)	ND(4.1)	ND
Angel shark (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	8.7	61	69.7
Sea raven (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.7)	ND(3.3)	ND
Common skate (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.6)	6.3	6.3
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(4.6)	4.5	4.5
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.7)	ND(3.5)	ND
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(4.2)	ND(4.6)	ND
Smooth dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.8)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(4.1)	ND(4.2)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(3.6)	6.0	6.0
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	Jan. 16, 2017	ND(2.8)	ND(3.5)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(4.3)	ND(3.4)	ND
Stingray (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.3)	5.2	5.2
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(4.3)	ND(3.7)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.7)	ND(3.9)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.6)	7.7	7.7
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(2.9)	ND(4.1)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.6)	14	14

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(4.0)	ND(4.4)	ND
Sea robin (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.4)	ND(3.5)	ND
Dasyatis matsubarae (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.8)	ND(3.6)	ND
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.2)	ND(3.7)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.8)	ND(3.7)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.4)	4.4	4.4
Pagrus major (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(4.4)	ND(4.0)	ND
Pacific cod (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(3.3)	ND(3.9)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	Jan. 13, 2017	ND(4.3)	ND(4.4)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.1)	3.4	3.4

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.1)	3.8	3.8
Andrea cuttlefish (whole)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.1)	ND(4.1)	ND
Striped jewfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.6)	ND(3.1)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.4)	ND(3.6)	ND
Lophius litilon (whole)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.4)	ND(3.8)	ND
Gnathophis nystromi nystromi (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.7)	ND(2.9)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.9)	ND(3.7)	ND
Pennahia argentata (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.2)	ND(4.0)	ND
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.4)	ND(3.9)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.6)	ND(3.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.0)	5.0	5.0
Sea robin (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.9)	ND(3.9)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.0)	ND(4.1)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.1)	4.0	4.0
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.8)	6.7	6.7
Pagrus major (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.5)	ND(3.8)	ND
Dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.2)	ND(2.9)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(4.8)	ND(3.0)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	Jan. 13, 2017	ND(3.7)	ND(4.0)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.7)	ND(3.6)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Angel shark (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	6.2	34	40.2
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.4)	ND(3.1)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.8)	11	11
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.4)	ND(3.8)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.2)	9.0	9.0
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.9)	ND(3.1)	ND
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.5)	ND(3.7)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.3)	4.6	4.6
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.7)	ND(3.7)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.6)	11	11

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <13/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pagrus major (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	Jan. 28, 2017	ND(3.2)	ND(4.5)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.9)	ND(4.5)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.8)	ND(3.2)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.8)	ND(4.1)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.9)	ND(4.5)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.9)	ND(4.2)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.1)	3.7	3.7
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(4.1)	4.5	4.5
Sea robin (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(4.0)	ND(4.2)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.2)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <14/14>
(excluding the port)

(Data summarized on February 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.2)	ND(3.7)	ND
Pagrus major (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(4.0)	ND(3.6)	ND
Common Octopus (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	Jan. 28, 2017	ND(3.8)	ND(3.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.