

**Nuclide Analysis Results of Fish and Shellfish**  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Spiny dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 2)	ND (3. 4)	ND
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 8)	ND (3. 0)	ND
Lepidotrigla microptera (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 9)	ND (3. 7)	ND
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (4. 1)	ND (3. 5)	ND
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 1)	5. 2	5. 2
Drumfish (whole)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (4. 0)	ND (4. 0)	ND
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 4)	ND (2. 9)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 8)	ND (3. 8)	ND
Smooth dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (4. 5)	ND (3. 6)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	May 11, 2017	ND (3. 7)	ND (3. 7)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Spiny dogfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 6)	ND (3. 8)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 1)	ND (3. 5)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (4. 4)	ND (3. 9)	ND
Lepidotrigla microptera (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 7)	ND (3. 7)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 6)	ND (3. 9)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 9)	9. 4	9. 4
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 3)	ND (3. 2)	ND
Ovalipes punctaus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 5)	ND (3. 0)	ND
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (4. 0)	ND (3. 7)	ND
Sea robin (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 3)	ND (3. 3)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Smooth dogfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 6)	ND (3. 6)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (4. 0)	ND (4. 2)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 11, 2017	ND (3. 7)	ND (3. 7)	ND
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 7)	ND (3. 6)	ND
Spiny dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (4. 0)	ND (3. 8)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 1)	ND (3. 2)	ND
Lepidotrigla microptera (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (4. 0)	ND (4. 1)	ND
Gray antimony(whole)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 5)	ND (4. 0)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 7)	12	12
Microstomus achne (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 6)	7. 2	7. 2

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Ovalipes punctatus (whole)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (4. 2)	ND (4. 0)	ND
Flatfish① (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (4. 0)	ND (3. 5)	ND
Flatfish② (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (4. 1)	ND (3. 5)	ND
Sea robin (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 8)	ND (3. 7)	ND
Pitted stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 7)	ND (3. 4)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 5)	ND (3. 9)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 7)	ND (3. 8)	ND
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (4. 0)	ND (3. 7)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 30, 2017	ND (3. 4)	ND (4. 7)	ND
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (4. 0)	ND (3. 7)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Gray antimony (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (4. 1)	ND (4. 1)	ND
Sea raven (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 8)	ND (3. 1)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 7)	6. 6	6. 6
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (4. 1)	7. 6	7. 6
Flatfish① (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 9)	5. 5	5. 5
Flatfish② (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 5)	ND (3. 9)	ND
Pitted stingray (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (4. 6)	ND (3. 6)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 9)	ND (4. 2)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 3)	ND (4. 2)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 30, 2017	ND (3. 5)	ND (3. 9)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Spiny dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (4. 2)	ND (3. 6)	ND
Sea raven (muscle)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (3. 6)	ND (3. 6)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (4. 3)	7. 7	7. 7
Drumfish (whole)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (4. 1)	ND (3. 7)	ND
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (4. 0)	4. 6	4. 6
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (3. 4)	ND (3. 6)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	May 17, 2017	ND (3. 7)	ND (4. 6)	ND
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (3. 2)	5. 9	5. 9
Spiny dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (3. 7)	ND (3. 8)	ND
Stone flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (4. 4)	ND (3. 6)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Fox jacopever (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (4. 0)	20	20
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (4. 5)	6. 1	6. 1
Sea bass (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (3. 1)	ND (4. 0)	ND
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (3. 1)	9. 3	9. 3
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (3. 9)	ND (3. 8)	ND
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 17, 2017	ND (4. 9)	7. 2	7. 2
Stone flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (3. 3)	ND (2. 9)	ND
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (4. 1)	7. 9	7. 9
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (3. 9)	7. 6	7. 6
Ovalipes punctaus (whole)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (4. 2)	ND (3. 9)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (4. 2)	ND (3. 8)	ND
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (4. 0)	ND (3. 3)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 9, 2017	ND (3. 4)	3. 8	3. 8
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 7)	ND (4. 0)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 5)	ND (3. 7)	ND
Lepidotrigla microptera (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (4. 5)	6. 5	6. 5
Gray antimony(whole)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 9)	ND (3. 9)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 6)	4. 6	4. 6
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 6)	ND (3. 4)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 4)	ND (3. 6)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.



Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 9)	ND (3. 4)	ND
Flatfish② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (3. 7)	ND (3. 5)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 19, 2017	ND (2. 8)	ND (3. 6)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 4)	4. 2	4. 2
Stone flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (2. 6)	ND (3. 7)	ND
Striped jewfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (4. 1)	ND (2. 8)	ND
Lepidotrigla microptera (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (4. 0)	ND (4. 0)	ND
Gray antimony(whole)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (2. 9)	ND (2. 8)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 7)	7. 0	7. 0
white croaker (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 7)	ND (3. 3)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/13>**  
**(excluding the port)**

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 6)	3. 5	3. 5
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (2. 6)	3. 6	3. 6
Flatfish① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (2. 9)	ND (3. 6)	ND
Flatfish② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (4. 6)	ND (3. 2)	ND
slime flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 6)	ND (3. 9)	ND
Japanese horse mackerel (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 3)	ND (4. 3)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 3)	ND (3. 8)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (4. 2)	ND (3. 2)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (4. 2)	5. 5	5. 5
red sea bream (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 8)	ND (2. 8)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/13>**  
**(excluding the port)**

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 9)	ND (4. 3)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 8)	ND (3. 2)	ND
Chestnut octopus(muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (4. 0)	ND (4. 3)	ND
willow flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 19, 2017	ND (3. 9)	ND (3. 7)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (3. 9)	4. 5	4. 5
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (4. 1)	ND (3. 4)	ND
Gray antimony(whole)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (4. 0)	ND (3. 8)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (3. 6)	ND (4. 5)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (3. 4)	3. 8	3. 8
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (4. 4)	ND (4. 2)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/13>**  
**(excluding the port)**

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (2. 7)	ND (3. 4)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (4. 2)	4. 0	4. 0
Sea robin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (3. 7)	ND (3. 4)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (3. 1)	ND (3. 7)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 8, 2017	ND (3. 3)	ND (3. 7)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 8)	ND (4. 1)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 3)	ND (3. 1)	ND
Lepidotrigla microptera (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 4)	ND (4. 3)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 7)	11	11
Sea bass (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 7)	4. 3	4. 3

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <13/13>  
(excluding the port)

(Data summarized on June 16)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 3)	ND (3. 5)	ND
Flatfish① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (4. 4)	ND (4. 4)	ND
Flatfish② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (4. 0)	6. 5	6. 5
Sea robin (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 0)	ND (3. 7)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 6)	ND (3. 2)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 7)	ND (3. 3)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (4. 1)	ND (3. 7)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (4. 1)	ND (3. 0)	ND
frog flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 8, 2017	ND (3. 3)	ND (3. 4)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.