

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stingray (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.9)	ND(3.5)	ND
Blue crab (whole)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(4.3)	ND(4.4)	ND
Japanese angel shark (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.9)	5.8	5.8
Gurnard (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.3)	ND(3.7)	ND
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.1)	ND(3.4)	ND
Drumfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.9)	ND(3.3)	ND
Slime flounder (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.7)	4.7	4.7
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.0)	ND(3.3)	ND
Species of hound shark (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.8)	ND(3.8)	ND
Chub mackerel (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2018	ND(3.9)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stingray (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(4.3)	ND(3.9)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(2.9)	ND(3.2)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.8)	ND(4.1)	ND
Gurnard (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.6)	ND(3.6)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.1)	4.8	4.8
Slime flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.9)	ND(3.6)	ND
Ovalipes punctatus (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.0)	ND(3.6)	ND
Flatfish ① (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(4.0)	ND(3.9)	ND
Flatfish ② (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.4)	ND(4.1)	ND
Sea robin (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.2)	ND(3.7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Pitted stingray (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.9)	ND(4.3)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.0)	ND(3.3)	ND
Chub mackerel (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.9)	ND(3.0)	ND
John Dory (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.0)	ND(4.0)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2018	ND(3.8)	ND(3.8)	ND
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.9)	20	20
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(2.9)	ND(3.9)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(4.2)	ND(4.3)	ND
Gurnard (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.6)	ND(3.5)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.4)	4.3	4.3

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Ovalipes punctatus (whole)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.5)	ND(4.2)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(4.0)	ND(3.9)	ND
Brown sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.2)	4.5	4.5
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.4)	ND(3.5)	ND
Chub mackerel (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 11, 2018	ND(3.8)	ND(3.6)	ND
Spiny dogfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.3)	ND(3.3)	ND
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.4)	ND(3.6)	ND
Gurnard (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.7)	ND(3.5)	ND
Black rockfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(4.0)	ND(4.1)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.3)	4.9	4.9

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sebastes cheni (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.9)	3.8	3.8
Slime flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(4.2)	ND(3.8)	ND
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.6)	ND(3.7)	ND
Pitted stingray (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.4)	ND(4.1)	ND
Marbled sole (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.4)	ND(3.6)	ND
Chub mackerel (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.1)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 11, 2018	ND(3.8)	ND(4.1)	ND
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.5)	ND(4.0)	ND
Stone flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.6)	ND(3.4)	ND
Gurnard (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.7)	ND(3.8)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Black sea bream (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.7)	ND(3.3)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(2.9)	ND(4.2)	ND
Drumfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(2.7)	ND(3.4)	ND
Slime flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.8)	6.8	6.8
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.5)	ND(3.0)	ND
Brown sole (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.0)	ND(4.2)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	May 15, 2018	ND(3.1)	3.9	3.9
Spiny dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(3.2)	ND(3.1)	ND
Blue crab (whole)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(3.7)	ND(4.4)	ND
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(4.0)	4.2	4.2

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Slime flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(3.9)	6.8	6.8
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(4.1)	ND(4.2)	ND
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(3.3)	ND(3.5)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 15, 2018	ND(3.1)	ND(3.5)	ND
Spiny dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(3.1)	ND(3.7)	ND
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(3.2)	3.6	3.6
Gurnard (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(4.2)	ND(3.4)	ND
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(4.2)	ND(3.5)	ND
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(3.0)	ND(4.1)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(3.7)	4.0	4.0

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Chub mackerel (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 11, 2018	ND(3.7)	ND(3.8)	ND
Spiny dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.4)	ND(3.3)	ND
Common skate (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(4.6)	ND(3.6)	ND
Slime flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.6)	8.9	8.9
Ovalipes punctatus (whole)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.9)	ND(3.1)	ND
Flatfish ① (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.6)	ND(4.3)	ND
Flatfish ② (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(4.1)	ND(3.5)	ND
Sea robin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.6)	ND(4.3)	ND
Brown sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.3)	ND(4.3)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(4.0)	ND(3.8)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 24, 2018	ND(3.1)	ND(3.9)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.5)	ND(4.0)	ND
Gurnard (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.5)	ND(4.1)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.5)	ND(3.2)	ND
Slime flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(2.9)	ND(3.5)	ND
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(4.2)	ND(3.7)	ND
Brown sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.6)	ND(3.5)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.4)	ND(3.4)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.5)	ND(3.6)	ND
Spearg squid (whole)	Around 15km Offshore of Odaka Ward (T-B1)	May 22, 2018	ND(3.7)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(3.9)	7.6	7.6
Striped jewfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(2.9)	ND(3.9)	ND
Gurnard (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(3.3)	ND(3.7)	ND
Gray antimony (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(3.8)	ND(3.5)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(4.0)	ND(3.6)	ND
Slime flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(3.6)	ND(3.1)	ND
Flatfish ① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(2.8)	ND(4.3)	ND
Flatfish ② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(4.5)	ND(4.2)	ND
Brown sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(4.1)	ND(3.5)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(3.2)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(2.5)	5.7	5.7
John Dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(4.1)	ND(4.1)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 22, 2018	ND(3.6)	ND(2.3)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.5)	ND(3.5)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.3)	ND(3.9)	ND
Gurnard (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.0)	ND(2.9)	ND
Gray antimony (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.3)	ND(3.5)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(4.1)	ND(4.0)	ND
Crimson sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(4.2)	ND(3.8)	ND
Slime flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.4)	ND(3.5)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <12/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish ① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.8)	ND(3.7)	ND
Flatfish ② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(4.1)	ND(3.9)	ND
Brown sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.6)	ND(3.6)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(2.9)	ND(3.3)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(3.5)	ND(3.3)	ND
Frog flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 21, 2018	ND(4.9)	3.7	3.7
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.8)	ND(3.8)	ND
Gurnard (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.6)	3.9	3.9
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.8)	5.8	5.8
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(4.1)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <13/13>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Slime flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.9)	ND(3.9)	ND
Flatfish ① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(4.3)	ND(3.3)	ND
Flatfish ② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.9)	ND(3.7)	ND
Brown sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.1)	ND(4.3)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(4.0)	ND(3.8)	ND
John Dory (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.6)	ND(4.2)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(3.3)	ND(3.2)	ND
Frog flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(4.2)	ND(4.1)	ND
Willow flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 21, 2018	ND(4.0)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.