

Nuclide analysis results of fish and shellfish
<The marine area where 20km of Fukushima Daiichi Nuclear Power Station>

(Data summarized on April 12)

Name of Sample (parts)	Place of Sampling	Day of Sampling	Radioactivity density (Bq/kg (raw)) (Half-life)		
			Cs-134 (approx. 2 years)	Cs-137 (approx. 30 years)	I-131 (approx. 8 days)
Ishikawasirauo (whole)	2km offshore of the Kido-gawa River	Mar 29, 2012	11	12	ND
Kounago (whole)	2km offshore of the Kido-gawa River	Mar 29, 2012	4.9	8.0	ND
Kounago (whole)	5km offshore of the Kido-gawa River	Mar 29, 2012	ND	ND	ND

* In the case the measurement is under the detection threshold, "ND" is marked.

Cs-134: approx. 3.6Bq/kg(raw)、Cs-137: approx. 4.1Bq/kg(raw)、I-131: approx. 11Bq/kg(raw).

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.

* Provisional Regulation Values (Until Mar. 31, 2012) Radioactive Iodine: 2000Bq/kg、Radioactive Cesium: 500Bq/kg.

* Standard limits for Radionuclides in Foods (Since Apr. 1, 2012) Sum of Cs-134 and Cs-137: 100Bq/kg.