

Correct

Nuclide Analysis Results of Fish and Shellfish
(The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 2/4 >

- List of Errata for the report as of June 8, 2012 -

(Data summarized on June 8)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	Total
Octopus (Enteroctopus) dofleini (Muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	May 21, 2012	ND	ND	ND
Roundnose flounder (Muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	May 21, 2012	25	32	57
Greenling (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	360	550	910
Schlegel's black rockfish (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	170	240	410
Common Skete (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	210	320	530
Drumfish (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	53	68	121
Microstomus achne (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	180	240	420
Flatfish (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	73	100	173
Dasyatis matsubarai (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	85	120	205
Barfin flounder (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	280	410	690

* When the measurement value is below the detection limit, "ND" is marked. The detection limits are as follows.

Cs-134: Approx. 3.7Bq/kg (Raw), Cs-137: Approx. 4.7Bq/kg (Raw).

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Original

Nuclide Analysis Results of Fish and Shellfish
(The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 2/4 >

(Data summarized on June 8)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	Total
Octopus (Enteroctopus) dofleini (Muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	May 21, 2012	ND	ND	ND
Roundnose flounder (Muscle)	Around 10km Offshore of Fukushima Daini NPS (T-B4)	May 21, 2012	25	32	57
Greenling (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	360	550	910
Schlegel's black rockfish (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	170	240	410
Common Skete (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	210	320	530
Pennahia argentata (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	53	68	121
Microstomus achne (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	180	240	420
Flatfish (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	73	100	173
Dasyatis matsubarai (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	85	120	205
Barfin flounder (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	280	410	690

* When the measurement value is below the detection limit, "ND" is marked. The detection limits are as follows.

Cs-134: Approx. 3.7Bq/kg (Raw), Cs-137: Approx. 4.7Bq/kg (Raw).

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Correct

Nuclide Analysis Results of Fish and Shellfish
(The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 3/4 >
- List of Errata for the report as of June 8, 2012 -

(Data summarized on June 8)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	Total
Marbled sole (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	100	150	250
Greenling (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	200	280	480
Northern dogfish (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	ND	ND	ND
Lophius litilon (Whole)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	5.4	11	16.4
Sea raven (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	270	400	670
Common Skete (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	360	480	840
Pennahia argentata (incl. Drumfish) (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	30	39	69
Sebastes cheni (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	620	970	1590
Sea bass (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	21	33	54
Microstomus achne (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	210	330	540

* When the measurement value is below the detection limit, "ND" is marked. The detection limits are as follows.

Cs-134: Approx. 3.6Bq/kg (Raw), Cs-137: Approx. 4.2Bq/kg (Raw).

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Original

Nuclide Analysis Results of Fish and Shellfish
(The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 3/4 >

(Data summarized on June 8)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	Total
Marbled sole (Muscle)	Around 3km offshore of Odaka Ward (T-S3)	May 25, 2012	100	150	250
Greenling (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	200	280	480
Northern dogfish (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	ND	ND	ND
Lophius litilon (Whole)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	5.4	11	16.4
Sea raven (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	270	400	670
Common Skete (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	360	480	840
Pennahia argentata (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	30	39	69
Sebastes cheni (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	620	970	1590
Sea bass (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	21	33	54
Microstomus achne (Muscle)	Around 3km Offshore of Fukushima Daiichi NPS (T-S4)	May 25, 2012	210	330	540

* When the measurement value is below the detection limit, "ND" is marked. The detection limits are as follows.

Cs-134: Approx. 3.6Bq/kg (Raw), Cs-137: Approx. 4.2Bq/kg (Raw).

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

* Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.