

Correct

Nuclide Analysis Results of Fish and Shellfish  
(The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 1/5 >  
- List of Errata for the report as of June 15, 2012 -

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling ( Place No. )	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	Total
Stingray (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	22	33	55
Flatfish (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	120	170	290
Sea bass (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	140	210	350
Banded dogfish (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	190	300	490
Drumfish(Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	50	74	124
Common Skete (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	220	310	530
Microstomus achne (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	200	290	490
Marbled sole (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	220	300	520
Lophius litilon (Whole)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	4.4	5.6	10
Blue crab (Whole)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	ND	7.1	7.1

\* When the measurement value is below the detection limit, "ND" is marked. The detection limits are as follows.

Cs-134: Approx. 4.1Bq/kg (Raw).

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

\* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

\* Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.

Original

Nuclide Analysis Results of Fish and Shellfish  
(The Ocean Area Within 20km Radius of Fukushima Daiichi NPS) < 1/5 >

(Data summarized on June 15)

Name of Sample (Region)	Place of Sampling ( Place No. )	Date of Sampling	Radioactivity Density[Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	Total
Stingray (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	22	33	55
Flatfish (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	120	170	290
Sea bass (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	140	210	350
Banded dogfish (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	190	300	490
Pennahia argentata (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	50	74	124
Common Skete (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	220	310	530
Microstomus achne (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	200	290	490
Marbled sole (Muscle)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	220	300	520
Lophius litilon (Whole)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	4.4	5.6	10
Blue crab (Whole)	Around 2km Offshore of Fukushima Daini NPS (T-S7)	May 30, 2012	ND	7.1	7.1

\* When the measurement value is below the detection limit, "ND" is marked. The detection limits are as follows.

Cs-134: Approx. 4.1Bq/kg (Raw).

As the detection limit may vary depending on the detectors and sample properties, there are cases where nuclides below the detection limit are detected.

\* Standard Value (after April 1, 2012) Cs-134+Cs-137: 100Bq/kg

\* Analyzed by Tokyo Electric Power Environmental Engineering Co., Inc.