

The result of nuclide analysis <samples taken by a surveying ship "Kaiyou" of the Japan Coast Guard >

1. Radioactive concentration in seawater

Place of collection	Time and date of collection	Radioactive concentration (Bq/ℓ)								
		Upper layer			Middle layer			Lower layer		
		I-131	Cs-134	Cs-137	I-131	Cs-134	Cs-137	I-131	Cs-134	Cs-137
①	8:06 May 5, 2011	ND	ND	ND	ND	ND	ND	ND	ND	ND
②	10:29 May 5, 2011	ND	ND	ND	ND	ND	ND	ND	ND	ND
③	13:08 May 5, 2011	ND	ND	ND	ND	ND	ND	ND	ND	ND
④	15:00 May 5, 2011	ND	ND	ND	ND	ND	ND	ND	ND	ND
⑤	16:51 May 5, 2011	ND	ND	ND	ND	ND	ND	ND	ND	ND

Note) Detection limits are approx. 6Bq/ℓ for I-131, approx. 14Bq/ℓ for Cs-134 and approx. 14Bq/ℓ for Cs-137.

Place of collection	Depth of collection	
	Middle layer	Lower layer
①	57m	143m
②	268m	561m
③	379m	792m
④	200m	421m
⑤	53m	130m