Results of Nuclide Analysis of Seawater <Coast>

Reference

Place of Sampling		_	Channel of 5-6u 5-6u discharge			Around South Discharge Channel of 1F (appox. 330m south of 1-4u Discharge Channel)				of 2F In Discharge In Discharge In Discharge In Discharge	of 2F		(Bq/L)
Time and Date of Sample Collection	, ,	2011/06/13 2011/06/13 10:20 14:15			2011/06 9:25			2011/06/13 14:00		2011/06/13 9:20		6/13 5	(the density limit in the water outside of surrounding
Detected Nuclides (Half-life)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (1)/2)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	①Density of Sample (Bq/L)	Scaling Factor (①/②)	monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)	ND	-	ND	-	ND	-	ND	-	ND	-	ND	-	40
Cs-134 (about 2 years)	21	0. 35	ND	ı	24	0. 40	34	0. 57	ND	-	ND	-	60
Cs-137 (about 30 years)	30	0. 33	16	0. 18	25	0. 28	35	0. 39	ND	-	ND	-	90

X Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L

X Data of other nuclides are under evaluation.

[💥] In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

Results of Nuclide Analysis of Seawater $\langle 0ffshore 1/3 \rangle$

Referemce

											(2 5 5 5 5		200 011 0/ 11/
Place of Sampling	15 km offshore of MinamiSouma City Upper layer		15 km offshore of MinamiSouma City Lower layer		15 km offshore of Ukedo-gawa Upper layer		15 km offshore of Ukedo-gawa Lower layer		15 km offshore of Fukushima Daiichi Upper layer		15 km offshore of Fukushima Daiichi Lower layer		② Density limit by the announcement of Reactor Regulation
Time and Date of Sample Collection		2011/06/13 Sampling suspended		2011/06/13 Sampling suspended		No planned sampling		No planned sampling		No planned sampling		ampling	(Bq/L) (the density limit
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (①/②)	of surrounding monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)													40
Cs-134 (about 2 years)													60
Cs-137 (about 30 years)													90

Place of Sampling	15 km offshore of 15 km offshore Fukushima Daini Fukushima Dai Upper layer Lower layer		Daini	15 km offshore of Iwasawa Shore Upper layer		15 km offshore of Iwasawa Shore Lower layer		15 km offshore of Hirono- machi Upper layer		15 km offshore of Hirono- machi Lower layer		② Density limit by the announcement of Reactor Regulation	
Time and Date of Sample Collection	No planned sampling		No planned sampling		2011/06/13 Sampling suspended		2011/06/13 Sampling suspended		2011/06/13 7:05		2011/06/13 7:05		(Bq/L) (the density limit in the water outside
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	of surrounding monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)									ND	-	ND	-	40
Cs-134 (about 2 years)									ND	ı	ND	-	60
Cs-137 (about 30 years)									ND	-	ND	-	90

X Density by the announcement of Reactor Regulation is stated with an amount converted from Bg/cm3 to Bg/L

X Data of other nuclides are under evaluation.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

In the case that the data is below measurable limit, "ND" is stated.
Detection limits of the three main nuclides are as follows: I-131: approx. 6Bq/L, Cs-134: approx. 14Bq/L, Cs-137: approx. 15Bq/L
However, detection limits differs depending on the detectors and samples types, and therefore may be detected, under figures below.

Results of Nuclide Analysis of Seawater $\langle 0ffshore 2/3 \rangle$

Referemce

Place of Sampling Time and Date of Sample Collection	15 km offshore of MinamiSouma City Upper layer 2011/06/13 Sampling suspended		City MinamiSouma City Lower layer 3 2011/06/13		15 km offshore of Ukedo-gawa Upper layer 2011/06/13 Sampling suspended		15 km offshore of Ukedo-gawa Lower layer 2011/06/13 Sampling suspended		15 km offshore of Fukushima Daiichi Upper layer 2011/06/13 7:40		15 km offshore of Fukushima Daiichi Lower layer 2011/06/13 7:40		② Density limit by the announcement of Reactor Regulation (Bq/L) (the density limit in the water outside
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm3)	Scaling Factor (①/②)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	of surrounding monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)									ND	-	ND	-	40
Cs-134 (about 2 years)									ND	-	ND	-	60
Cs-137 (about 30 years)									ND	-	ND	-	90

Place of Sampling Time and Date of Sample Collection	15 km offshore of Fukushima Daini Upper layer 2011/06/13 Sampling suspended		15 km offshore of Fukushima Daini Lower layer 2011/06/13 Sampling suspended		15 km offshore of Iwasawa Shore Upper layer 2011/06/13 7:40		15 km offshore of Iwasawa Shore Lower layer 2011/06/13 7:40						② Density limit by the announcement of Reactor Regulation (Bq/L) (the density limit
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (①/②)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	in the water outside of surrounding monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)					ND	-	ND	-					40
Cs-134 (about 2 years)					ND	-	ND	-					60
Cs-137 (about 30 years)					ND	-	ND	_					90

X Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L

X Data of other nuclides are under evaluation.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

In the case that the data is below measurable limit, "ND" is stated.
Detection limits of the three main nuclides are as follows: I-131: approx. 6Bq/L, Cs-134: approx. 14Bq/L, Cs-137: approx. 15Bq/L
However, detection limits differs depending on the detectors and samples types, and therefore may be detected, under figures below.

Results of Nuclide Analysis of Seawater <0ffshore 3/3>

Referemce

											(20.00.		200 011 0/ 11/
Place of Sampling	15 km offshore of MinamiSouma City Upper layer		15 km offshore of MinamiSouma City Lower layer		15 km offshore of Ukedo-gawa Upper layer		Ukedo-ga	15 km offshore of Ukedo-gawa Lower layer		15 km offshore of Fukushima Daiichi Upper layer		ore of Daiichi yer	2 Density limit by the announcement of Reactor Regulation
Time and Date of Sample Collection	2011/06 6:00		2011/06, 6:00		2011/06, 5:40		2011/06 5:40		2011/06, 5:45	/13	2011/06/13 5:45		(Bq/L) (the density limit in the water outside
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	of surrounding monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)	ND	_	ND	-	ND	-	ND	-	ND	-	ND	_	40
Cs-134 (about 2 years)	ND	-	ND	-	4. 5	0. 08	ND	-	ND	-	ND	-	60
Cs-137 (about 30 years)	ND	-	ND	-	ND	-	ND	-	ND	-	ND	-	90

Place of Sampling			Shore	15 km offshore of Iwasawa Shore Upper layer		15 km offshore of Iwasawa Shore Lower layer		of Hirono- i yer	15 km offshore of Hirono- machi Lower layer		2 Density limit by the announcement of Reactor Regulation		
Time and Date of Sample Collection	2011/06, 6:00		2011/06, 6:00		2011/06 5:30		2011/06 5:30		2011/06 5:20		2011/06 5:20		(Bq/L) (the density limit in the water outside
Detected Nuclides (Half-life)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	①Density of Sample (Bq/cm3)	Scaling Factor (1)/2)	of surrounding monitored areas in the section 6 of the appendix 2)
I-131 (about 8 days)	ND	-	ND	-	ND	-	ND	-	ND	-	ND	-	40
Cs-134 (about 2 years)	ND	_	ND	_	ND	-	ND	_	ND	-	ND	_	60
Cs-137 (about 30 years)	ND	-	ND	-	ND	_	ND	-	ND	-	ND	-	90

X Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L

X Data of other nuclides are under evaluation.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

In the case that the data is below measurable limit, "ND" is stated.
Detection limits of the three main nuclides are as follows: I-131: approx. 6Bq/L, Cs-134: approx. 14Bq/L, Cs-137: approx. 15Bq/L
However, detection limits differs depending on the detectors and samples types, and therefore may be detected, under figures below.