

Nuclide analysis results of ocean soil

Reference

(Data summarized on September 13)

| Place of Sampling | North of Discharge Channel of 5-6u of 1F (approx. 30m north of 5-6u discharge channel) | Around South Discharge Channel of 1F (approx. 330m south of 1-4u Discharge Channel) | Around North Discharge Channel of 2F (Around 3,4u Discharge Channel) (approx. 10 km from 1F) | Around Iwasawa Shore of 2F (approx. 7 km south of 1,2u Discharge Channel) (approx. 16 km from 1F) |
|-------------------------------|--|--|---|--|
| Time of Sampling | 10:45 Sep 12 2011 | 2011 Sep 12 (Not sampled) | 2011 Sep 12 (Not sampled) | 2011 Sep 12 (Not sampled) |
| Detected Nuclides (Half-life) | Density of sample (Bq/kg) | | | |
| I-131 (about 8 days) | ND | - | - | - |
| Cs-134 (about 2 years) | 2,000 | - | - | - |
| Cs-137 (about 30 years) | 2,300 | - | - | - |

* Data of other nuclides are under evaluation.

* "ND" means the sampled data is below measurable limit. I-131: approx. 14Bq/kg)

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.