

Nuclide analysis results of ocean soil (Additional)

Reference

(Data summarized on September 30)

Place of Sampling	Offshore of Soma 10km	Offshore of Kashima 3km	Offshore of Kashima 10km	Offshore of Kashima 15km	Offshore of Mano River 3km	Offshore of Mano River 5km	Offshore of Mano River 10km	Offshore of Mano River 15km
Time of Sampling	2011/9/27 5:55 AM	2011/9/28 6:32 AM	2011/9/27 6:13 AM	2011/9/26 6:15 AM	2011/9/28 6:56 AM	2011/9/28 7:21 AM	2011/9/27 6:31 AM	2011/9/26 6:41 AM
Detected Nuclides (Half-life)	Density of sample (Bq/kg)							
I-131 (about 8 days)	ND	ND	ND	ND	ND	ND	ND	ND
Cs-134 (about 2 years)	14	110	54	22	47	27	22	22
Cs-137 (about 30 years)	17	130	56	27	61	33	27	25

Place of Sampling	Offshore of Yotsukura 3km	Offshore of Yotsukura 8km	Offshore of Yotsukura 15km	Offshore of Natsui River 8km	Offshore of Natsui River 15km	Offshore of Toyoma 5km	Offshore of Ena 5km	Offshore of Onahama East 5 km
Time of Sampling	2011/9/28 5:10 AM	2011/9/28 5:55 AM	2011/9/27 5:40 AM	2011/9/28 6:25 AM	2011/9/27 6:20 AM	2011/9/26 6:42 AM	2011/9/26 6:12 AM	2011/9/26 5:50 AM
Detected Nuclides (Half-life)	Density of sample (Bq/kg)							
I-131 (about 8 days)	ND	ND	ND	ND	ND	ND	ND	ND
Cs-134 (about 2 years)	190	250	56	130	20	64	320	1,100
Cs-137 (about 30 years)	230	290	63	150	24	83	370	1,300

* Data of other nuclides are under evaluation.

* "ND" means the sampled data is below measurable limit. I-131: approx. 11Bq/kg

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.