Nuclide Analysis Results of Seawater < Coast>

Reference

(Data summarized on December 20)

| Place of Sampling | North of Discharge Channel of 5-6u of 1F (approx. 30m north of 5-6u discharge channel) | | Around South Discharge Channel of 1F (appox. 330m south of 1-4u Discharge Channel) | | Around North Discharge Channel of 2F (Around 3,4u Discharge Channel) (approx. 10 km from 1F) | | Around Iwasawa Shore of 2F (appox. 7 km south of 1,2u Discharge Channel) (appox. 16 km from 1F) | | Danaite dissit beet the |
|----------------------------------|---|----------------------------|--|----------------------------|---|----------------------------|---|----------------------------|---|
| Time of Sampling | Dec 19, 2011 08:35 am | | Dec 19, 2011 08:15 am | | Dec 19, 2011 08:20 am | | 07:55 am | | |
| Detected Nuclides (Half-life) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | Density of Sample (Bq/L) | Scaling Factor (/) | areas in the section 6 of the appendix 2) |
| I-131 (about 8 days) | ND | - | ND | - | ND | - | ND | - | 40 |
| Cs-134 (about 2 years) | 2.7 | 0.05 | 1.1 | 0.02 | 0.91 | 0.02 | ND | - | 60 |
| Cs-137 (about 30 years) | 2.7 | 0.03 | 1.7 | 0.02 | 1.1 | 0.01 | ND | - | 90 |

^{*} Density by the announcement of Reactor Regulation is stated with an amount converted from Bq/cm3 to Bq/L.

Please note that these nuclides are sometimes detected even when they are below the limits, contingent on the detector or samples.

^{*} Data of other nuclides are under evaluation.

^{*} In the case that two or more kinds of nuclides exist, sum of each scaling factor to the density limit is compared with 1.

^{* &}quot;ND" means the sampled data is below measurable limit.

I-131: approx. 0.73Bq/L, Cs-134: approx. 0.86Bq/L, Cs-137: approx. 0.98Bq/L