

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Japanese angel shark (muscle)	Around 1km Offshore of Ota River (T-S1)	December 6, 2018	ND(4.1)	ND(3.6)	ND
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	December 6, 2018	ND(3.3)	ND(3.8)	ND
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	December 6, 2018	ND(3.7)	4.8	4.8
Japanese amberjack (muscle)	Around 1km Offshore of Ota River (T-S1)	December 6, 2018	ND(3.4)	ND(3.8)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	December 6, 2018	ND(3.7)	ND(4.0)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.7)	ND(3.2)	ND
Lepidotrigla microptena (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.3)	ND(4.1)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.8)	ND(3.6)	ND
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.0)	ND(3.5)	ND
Searobin (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.9)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.1)	ND(3.4)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	December 6, 2018	ND(3.6)	ND(3.9)	ND
Stingray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(3.9)	ND(4.5)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(3.6)	ND(3.4)	ND
Common skate (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(4.1)	ND(3.6)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(3.5)	ND(3.7)	ND
Searobin (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(2.8)	ND(3.8)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(3.9)	ND(3.1)	ND
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	December 14, 2018	ND(3.7)	ND(3.5)	ND
Stingray (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(4.1)	ND(3.7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Yellow goosfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(4.2)	ND(4.2)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(4.1)	ND(4.0)	ND
Flatfish ① (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(3.8)	ND(3.2)	ND
Flatfish ② (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(4.1)	ND(3.9)	ND
Searobin (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(4.0)	ND(3.3)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	December 14, 2018	ND(3.8)	ND(3.5)	ND
Black sea bream (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(4.1)	ND(3.8)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.4)	ND(3.9)	ND
Banded houndshark (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.5)	ND(3.3)	ND
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.7)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(4.0)	ND(4.2)	ND
Searobin (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.4)	ND(3.8)	ND
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.0)	ND(3.4)	ND
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.5)	ND(3.9)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	December 4, 2018	ND(3.7)	ND(3.6)	ND
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	December 4, 2018	ND(3.0)	ND(4.2)	ND
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	December 4, 2018	ND(4.0)	7.7	7.7
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	December 4, 2018	ND(4.0)	ND(3.8)	ND
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	December 4, 2018	ND(4.0)	ND(3.4)	ND
Stingray (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.9)	4.2	4.2

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.0)	ND(3.7)	ND
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(4.0)	3.9	3.9
Flatfish ① (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.4)	ND(3.6)	ND
Flatfish ② (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(4.0)	ND(4.0)	ND
Searobin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(4.3)	ND(3.9)	ND
Smooth dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.3)	ND(4.0)	ND
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.6)	ND(4.0)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.5)	3.8	3.8
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	December 20, 2018	ND(3.3)	ND(3.5)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.9)	ND(4.0)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Lepidotrigla microptena (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.2)	ND(3.8)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.9)	ND(4.1)	ND
Globefish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.3)	ND(4.3)	ND
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.5)	ND(3.0)	ND
Crimson sea bream (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.7)	ND(3.5)	ND
Flatfish ① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.4)	ND(3.6)	ND
Flatfish ② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.7)	ND(4.3)	ND
Searobin (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.5)	ND(3.2)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.5)	ND(3.9)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(4.1)	ND(4.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Red sea bream (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.7)	ND(3.6)	ND
John dory (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(4.5)	ND(3.2)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	December 11, 2018	ND(3.0)	ND(3.9)	ND
Stingray (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.3)	ND(3.9)	ND
Stone flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(4.4)	ND(4.5)	ND
Lepidotrigla microptena (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(4.1)	ND(3.5)	ND
Yellow goosefish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.6)	ND(3.8)	ND
Conger eel (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(2.6)	ND(3.8)	ND
Common skate (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.9)	ND(3.8)	ND
Globefish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.1)	ND(4.1)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Sea bass (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(4.2)	ND(3.7)	ND
Crimson sea bream (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.6)	ND(3.4)	ND
Flatfish ① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.9)	ND(3.0)	ND
Flatfish ② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(4.4)	ND(2.8)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.0)	3.0	3.0
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.4)	ND(3.4)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.6)	ND(3.5)	ND
John dory (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.9)	ND(4.0)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.7)	ND(4.0)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	December 11, 2018	ND(3.3)	ND(3.6)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(4.4)	ND(3.6)	ND
Lepidotrigla microptena (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.4)	3.9	3.9
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.9)	ND(3.3)	ND
Globefish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.0)	ND(4.2)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.9)	ND(3.4)	ND
Crimson sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.9)	ND(3.2)	ND
Flatfish ① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(4.0)	ND(4.0)	ND
Flatfish ② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(4.0)	ND(3.7)	ND
Searobin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.1)	ND(3.6)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.8)	ND(3.6)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(4.0)	ND(3.7)	ND
Red sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(4.6)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(4.2)	ND(4.4)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	December 17, 2018	ND(3.9)	ND(4.1)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.9)	ND(3.8)	ND
Lepidotrigla microptena (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.0)	ND(4.2)	ND
Common skete (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.5)	ND(3.4)	ND
Globefish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(4.2)	ND(3.2)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.8)	ND(3.4)	ND
Flatfish ① (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.8)	ND(4.3)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish ② (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(4.7)	ND(3.4)	ND
Searobin (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.9)	ND(3.5)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.3)	ND(3.5)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.7)	ND(3.6)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.2)	ND(2.9)	ND
Red sea bream (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.6)	ND(3.6)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.8)	ND(3.5)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	December 17, 2018	ND(3.7)	ND(3.3)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.