

### Nuclide Analysis Results of Fish <Port Area of Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Black sea bream (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	December 3, 2018	ND(5.0)	22	22
Greenling (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	December 17, 2018	ND(4.7)	23	23
Schlegel's black rockfish (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	December 19, 2018	11	180	191
Sea bass (muscle) No.1	Port area of Fukushima Daiichi NPS (Near port entrance)	December 19, 2018	ND(8.0)	37	37
Drumfish (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	December 22, 2018	ND(6.4)	25	25
Sea bass (muscle) No.2	Port area of Fukushima Daiichi NPS (Near port entrance)	December 23, 2018	ND(4.1)	ND(4.2)	ND
Flatfish (muscle)	Port area of Fukushima Daiichi NPS (Center in open channel)	December 18, 2018	55	660	715
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS (Center in open channel)	December 18, 2018	59	600	659
Sebastes cheni (muscle)	Port area of Fukushima Daiichi NPS (North side of east breakwater)	December 26, 2018	44	550	594

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.