Nuclide Analysis Results of Fish < Port Area of Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS (North breakwater)	April 3, 2019	ND(4.7)	40	40
Flatfish (muscle)	Port area of Fukushima Daiichi NPS (South breakwater)	April 26, 2019	ND(4.9)	20	20
Temminck's surfperch (muscle)	Port area of Fukushima Daiichi NPS (North side in open channel)	April 3, 2019	ND(8.2)	39	39
Gray mullet (muscle)	Port area of Fukushima Daiichi NPS (North side in open channel)	April 19, 2019	13	160	173
Sea bass (muscle)	Port area of Fukushima Daiichi NPS (South side in open channel)	April 3, 2019	ND(4.7)	6.8	6.8

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.