

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(4.2)	ND(3.9)	ND
Lepidotrigla microptena (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(4.2)	ND(3.3)	ND
Yellow goosefish (whole)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.9)	ND(3.8)	ND
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.7)	ND(3.1)	ND
Black sea bream (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.1)	ND(3.7)	ND
Microstomus achne (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.0)	ND(3.5)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.8)	ND(3.5)	ND
Smooth dogfish (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.4)	ND(4.1)	ND
Marbled sole (muscle)	Around 1km Offshore of Ota River (T-S1)	April 18, 2019	ND(3.8)	ND(3.5)	ND
Stone flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.7)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.7)	ND(3.8)	ND
Lepidotrigla microptena (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.9)	ND(3.7)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(4.0)	ND(3.5)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.8)	ND(3.8)	ND
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.1)	ND(3.7)	ND
Flatfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.9)	ND(3.2)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.5)	ND(3.3)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	April 18, 2019	ND(3.3)	ND(3.6)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(3.5)	ND(3.8)	ND
Lepidotrigla microptena (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(3.8)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Schlegel's black rockfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(3.3)	ND(3.7)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(4.0)	ND(3.5)	ND
Sea bass (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(3.7)	ND(3.6)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(4.0)	ND(3.3)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(3.0)	5.0	5.0
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(4.2)	ND(3.0)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	April 4, 2019	ND(4.2)	3.0	3.0
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.9)	ND(3.2)	ND
Lepidotrigla microptena (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.7)	ND(3.4)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.9)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.9)	ND(3.3)	ND
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(4.1)	ND(4.0)	ND
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.5)	ND(3.5)	ND
Searobin (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.6)	ND(4.1)	ND
Smooth dogfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.4)	ND(4.0)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	April 4, 2019	ND(3.8)	ND(3.0)	ND
Schlegel's black rockfish (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(3.5)	ND(3.6)	ND
Black sea bream (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(4.0)	ND(4.1)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(3.7)	ND(3.2)	ND
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(3.9)	ND(3.6)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Searobin (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(4.0)	3.9	3.9
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(3.5)	ND(3.9)	ND
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(3.7)	ND(3.5)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	April 9, 2019	ND(3.0)	ND(3.9)	ND
Stone flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	April 9, 2019	ND(3.8)	ND(4.2)	ND
Schlegel's black rockfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	April 9, 2019	ND(3.9)	ND(3.6)	ND
Common skete (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	April 9, 2019	ND(3.4)	7.2	7.2
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	April 9, 2019	ND(3.9)	ND(3.4)	ND
Marbled sole (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	April 9, 2019	ND(3.7)	5.8	5.8
Greenling (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(4.3)	ND(3.7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Lepidotrigla microptena (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.9)	ND(3.4)	ND
Schlegel's black rockfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.6)	ND(3.2)	ND
Sea raven (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.6)	ND(4.3)	ND
Common skete (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.4)	ND(3.4)	ND
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.4)	4.0	4.0
Flatfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.8)	ND(3.2)	ND
Smooth dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.8)	ND(4.0)	ND
Littlemouth flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.4)	ND(3.7)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.6)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	April 10, 2019	ND(3.7)	ND(3.4)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.6)	ND(3.3)	ND
Black scraper (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(2.9)	ND(3.7)	ND
Lepidotrigla microptena (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.5)	ND(3.6)	ND
Yellow gosefish (whole)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.8)	ND(3.6)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(4.0)	ND(4.2)	ND
Sea bass (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.9)	ND(3.9)	ND
Pointhead flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.8)	ND(3.6)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.7)	ND(3.5)	ND
Flatfish ① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(4.1)	ND(3.5)	ND
Flatfish ② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.8)	ND(4.1)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.2)	3.8	3.8
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.6)	ND(3.9)	ND
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	April 16, 2019	ND(3.2)	ND(3.5)	ND
Greenling (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.6)	ND(3.6)	ND
Lepidotrigla microptena (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.7)	ND(4.1)	ND
Yellow gosefish (whole)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.5)	ND(4.1)	ND
Common skete (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.9)	ND(3.8)	ND
Pointhead flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(4.1)	ND(3.5)	ND
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.3)	ND(3.9)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(4.0)	ND(3.6)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.4)	ND(3.7)	ND
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.4)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.3)	ND(3.4)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	April 16, 2019	ND(3.7)	ND(3.4)	ND
Lepidotrigla microptena (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.8)	ND(3.7)	ND
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.5)	ND(3.3)	ND
Sea bass (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.5)	ND(3.6)	ND
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.2)	ND(3.6)	ND
Crimson sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.7)	ND(3.7)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(4.0)	ND(3.7)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish ① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.8)	ND(3.8)	ND
Flatfish ② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.6)	ND(3.4)	ND
Searobin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(4.6)	ND(3.8)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.6)	ND(3.3)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.0)	ND(3.8)	ND
Chub mackerel (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.3)	ND(3.4)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	April 2, 2019	ND(3.7)	ND(3.3)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.7)	ND(4.2)	ND
Lepidotrigla microptena (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.2)	ND(3.7)	ND
Yellow goosefish (whole)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.5)	ND(3.9)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/11>
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.1)	ND(3.8)	ND
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(2.8)	ND(3.3)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(4.3)	ND(3.4)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.0)	ND(3.5)	ND
Searobin (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.9)	ND(3.4)	ND
Smooth dogfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.6)	ND(3.2)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.6)	ND(3.5)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.8)	ND(3.9)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(3.2)	ND(3.6)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	April 2, 2019	ND(2.7)	ND(3.2)	ND

*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

*Analyzed by: Tokyo Power Technology Ltd.