Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Gizzard shad (muscle)	Port area of Fukushima Daiichi NPS (Near port entrance)	August 5, 2019	ND(5.0)	6.5	6.5
Flatfish (muscle) No.1	Port area of Fukushima Daiichi NPS (Near port entrance)	August 5, 2019	ND(3.1)	ND(3.1)	ND
Flatfish (muscle) No.2	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	August 1, 2019	ND(2.8)	3.8	3.8
Flatfish (muscle) No.3	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	August 29, 2019	ND(2.7)	20	20
Spotted halibut (muscle)	Port area of Fukushima Daiichi NPS (South side of the east breakwater)	August 29, 2019	ND(2.7)	59	59

^{*}ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

^{*}Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.