

**Nuclide Analysis Results of Fish and Shellfish**  
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <1/8>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	October 3, 2019	ND(3.2)	ND(4.4)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	October 3, 2019	ND(4.0)	ND(3.7)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	October 3, 2019	ND(3.9)	ND(3.0)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	October 3, 2019	ND(3.6)	ND(3.6)	ND
Blue crab (whole)	Around 3km Offshore of Ukedo River (T-S3)	October 10, 2019	ND(3.5)	ND(3.5)	ND
Japanese angel shark (muscle)	Around 3km Offshore of Ukedo River (T-S3)	October 10, 2019	ND(3.5)	ND(3.3)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	October 10, 2019	ND(3.7)	ND(3.8)	ND
Japanese eagle ray (muscle)	Around 3km Offshore of Ukedo River (T-S3)	October 10, 2019	ND(3.7)	ND(3.7)	ND
Searobin (muscle)	Around 3km Offshore of Ukedo River (T-S3)	October 10, 2019	ND(3.8)	ND(3.6)	ND
Red sea bream (muscle)	Around 3km Offshore of Ukedo River (T-S3)	October 10, 2019	ND(3.2)	ND(3.2)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <2/8>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Japanese angel shark (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	October 10, 2019	ND(4.0)	ND(3.4)	ND
Common skete (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	October 10, 2019	ND(3.2)	ND(3.8)	ND
Flatfish (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	October 10, 2019	ND(3.5)	ND(3.4)	ND
Searobin (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	October 10, 2019	ND(3.9)	ND(3.6)	ND
Stone flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(3.7)	ND(3.7)	ND
Blue crab (whole)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(3.5)	19	19
Japanese angel shark (muscle)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(4.0)	ND(3.4)	ND
Black sea bream (muscle)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(3.8)	ND(3.1)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(3.1)	ND(4.0)	ND
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(4.3)	ND(3.8)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <3/8>**  
**(excluding the port)**

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Smooth dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	October 31, 2019	ND(3.3)	ND(4.4)	ND
Sea raven (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	October 31, 2019	ND(3.3)	ND(3.2)	ND
Common skate (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	October 31, 2019	ND(4.4)	3.9	3.9
Sea bass (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	October 31, 2019	ND(4.0)	ND(4.1)	ND
Banded houndshark (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	October 31, 2019	ND(4.3)	ND(3.7)	ND
Smooth dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	October 31, 2019	ND(4.0)	ND(4.1)	ND
Stone flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(4.6)	ND(3.4)	ND
Blue crab (whole)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(4.7)	6.0	6.0
Japanese angel shark (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(3.6)	14	14
Sea raven (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(3.8)	ND(4.1)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <4/8>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
White croaker (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(3.4)	ND(3.4)	ND
Tiger puffer (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(3.6)	4.2	4.2
Flatfish ① (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(4.1)	ND(3.8)	ND
Flatfish ② (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(4.0)	ND(3.4)	ND
Searobin (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(3.8)	ND(3.9)	ND
Smooth dogfish (muscle)	Around 4km Offshore of Kumagawa (T-S8)	October 31, 2019	ND(3.8)	ND(4.0)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.9)	ND(3.7)	ND
Lepidotrigla microptena (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.3)	ND(3.3)	ND
Common skete (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(4.0)	ND(3.5)	ND
White croaker (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.1)	ND(3.7)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <5/8>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Crimson sea bream (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.7)	ND(4.0)	ND
Flatfish ① (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.8)	3.8	3.8
Flatfish ② (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(4.1)	ND(2.5)	ND
Searobin (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.3)	ND(3.8)	ND
Smooth dogfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(4.2)	ND(4.3)	ND
Marbled sole (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.6)	ND(3.7)	ND
Red sea bream (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.7)	ND(2.8)	ND
John dory (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.4)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.1)	ND(4.0)	ND
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	October 29, 2019	ND(3.0)	ND(3.1)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <6/8>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.2)	ND(3.5)	ND
White croaker (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(4.3)	ND(4.0)	ND
Flatfish ① (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.9)	ND(4.1)	ND
Flatfish ② (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.7)	ND(4.0)	ND
Common Japanese conger (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.2)	ND(3.0)	ND
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.1)	ND(3.3)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.1)	ND(3.9)	ND
Red sea bream (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.7)	ND(4.0)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(4.3)	ND(4.3)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	October 29, 2019	ND(3.9)	ND(3.4)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <7/8>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Globefish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	October 21, 2019	ND(4.1)	ND(4.3)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	October 21, 2019	ND(3.1)	ND(3.2)	ND
Searobin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	October 21, 2019	ND(3.3)	ND(3.4)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	October 21, 2019	ND(3.5)	ND(3.8)	ND
Red sea bream (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	October 21, 2019	ND(3.4)	ND(3.1)	ND
John dory (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	October 21, 2019	ND(2.9)	ND(3.1)	ND
Japanese angel shark (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	October 21, 2019	ND(3.6)	ND(3.5)	ND
Globefish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	October 21, 2019	ND(3.3)	ND(3.2)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	October 21, 2019	ND(3.1)	ND(3.7)	ND
Red sea bream (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	October 21, 2019	ND(3.4)	ND(4.1)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
 (Sampled from the Ocean Area within a 20km Radius of the Fukushima Daiichi Nuclear Power Station) <8/8>  
 (excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
John dory (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	October 21, 2019	ND(4.0)	ND(3.6)	ND
Carcharhinus (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	October 21, 2019	ND(3.1)	ND(3.9)	ND

\*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

\*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.

\*Analysis was conducted by Tokyo Power Technology Ltd.