

Nuclide Analysis Results of Fish <Sampled from the Port Area of the Fukushima Daiichi Nuclear Power Station>

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Port area of Fukushima Daiichi NPS (Near north breakwater)	October 10, 2019	ND(2.3)	26	26
Spottedtail morwong (muscle)	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	October 18, 2019	ND(2.7)	8.1	8.1
Sebastes cheni (muscle)	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	October 21, 2019	ND(3.0)	8.8	8.8
Flatfish (muscle)	Port area of Fukushima Daiichi NPS (North side of the east breakwater)	October 24, 2019	2.8	68	70.8
Spotted halibut (muscle)	Port area of Fukushima Daiichi NPS (South side of the east breakwater)	October 1, 2019	3.5	41	44.5

*ND indicates that a value is less than detection limits of radioactivity concentration. The detection limit for each nuclide is shown in parenthesis.

*Reference value (on and after April 1, 2012) Sum of radioactivity concentrations for Cs-134 and Cs-137: 100Bq/kg.